arXiv:0809.1262v1 [math.DS] 8 Sep 2008

COMBINATORICS AND TOPOLOGY OF STRAIGHTENING MAPS I : COMPACTNESS AND BIJECTIVITY

HIROYUKI INOU AND JAN KIWI

Abstract. We study the parameter space structure of degree $d \geq 3$ one complex variable polynomials as dynamical systems acting on $\mathbb{C}$. We introduce and study *straightening maps*. These maps are a natural higher degree generalization of the ones introduced by Douady and Hubbard to prove the existence of small copies of the Mandelbrot set inside itself. We establish that straightening maps are always injective and that their image contains all the corresponding hyperbolic systems. Also, we characterize straightening maps with compact domain. Moreover, we give two classes of bijective straightening maps. The first produces an infinite collection of embedded copies of the $(d-1)$-fold product of the Mandelbrot set in the connectedness locus of degree $d \geq 3$. The second produces an infinite collection of full families of quadratic connected filled Julia sets in the cubic connectedness locus, such that each filled Julia set is quasiconformally embedded.

Contents

Introduction 2
1. Outline of results 4
1.1. Combinatorially renormalizable polynomials 5
1.2. Mapping schemata 5
1.3. Universal polynomial model space 5
1.4. Polynomial-like maps over schemata 6
1.5. External markings 6
1.6. Standard external marking of polynomial maps in Poly(T) 6
1.7. Straightening of polynomial-like maps over reduced schemata 7
1.8. Renormalizable polynomials 8
1.9. Internal angle systems and induced external markings 9
1.10. Straightening map and injectivity 9
1.11. Onto hyperbolic maps 10
1.12. Compactness 10
1.13. Cubic reduced mapping schemata 11
1.14. Capture and Disjoint type straightenings 12
2. Polynomial-like maps 13
2.1. Proof of the straightening theorem 13
3. Laminations 14
3.1. Laminations 14
3.2. Invariant rational laminations: unlinked classes, fibers and critical elements 15
3.3. Renormalizations 20
3.4. Proof of Proposition 3.7 21

The second author was partially supported by Research Network on Low Dimensional Systems, PBCT/CONICYT, Chile.

4. Yoccoz puzzles 22
4.1. Combinatorial Yoccoz Puzzle 22
4.2. Yoccoz Puzzle 23
4.3. From combinatorial puzzles to laminations 25
5. Injectivity of straightening maps 27
5.1. Proof of Theorem B 27
5.2. Proof of the key lemma 29
6. Combinatorial tuning 34
6.1. Combinatorial straightening and renormalized lamination 35
6.2. Combinatorial Tuning 35
6.3. Renormalizable set 38
6.4. Proof of Theorem 6.3 39
6.5. Proof of Theorem C 40
7. Compactness 40
8. Surjectivity of straightening maps 44
8.1. Preliminaries 44
8.2. Surjectivity of straightening: primitive disjoint type 45
8.3. Surjectivity of straightening: cubic primitive capture type 46
8.4. Complex submanifolds and quasiconformality of straightening 47
8.4.1. Proof of Theorem E 48
8.4.2. Capture case 48
References 49

INTRODUCTION

The Mandelbrot set consists of all parameters $c \in \mathbb{C}$ for which the filled Julia set $K(z^2+c)$ of the quadratic polynomial $z^2 + c$ is connected. In any reasonable picture of the Mandelbrot set $\mathcal{M}$ one immediately observes the presence of small copies of it contained in itself. Douady and Hubbard [DH2] gave a mathematical proof of this fact. They established the existence of homeomorphisms χ from carefully chosen subsets M' of the Mandelbrot set onto the whole Mandelbrot set $\mathcal{M}$. These homeomorphisms $\chi : M' \to \mathcal{M}$ are fundamental examples of what are usually called "straightening maps". The degree $d \geq 3$ analogue of the Mandelbrot set is called the connectedness locus $\mathcal{C}(d)$ of degree d. In this paper we propose a definition of straightening maps in the context of higher degree polynomial dynamics. This involves introducing appropriate domains as well as the corresponding target sets and the maps themselves. In contrast with quadratic polynomials, one encounters a large diversity of target sets. Therefore, a greater variety of structures replicate all over the corresponding connectedness loci. In particular, for cubic polynomials, the target sets include the cubic connectedness locus $\mathcal{C}(3)$, the connectedness locus of biquadratic polynomials $\{(a, b) \in \mathbb{C}^2;\ K((z^2 + a)^2 + b)) \text{ is connected}\}$, the full family of quadratic connected filled Julia sets $\mathcal{MK} = \{(c, z) \in \mathbb{C}^2;\ c \in \mathcal{M}, z \in K(z^2 + c)\}$, and the cross product of the Mandelbrot set with itself $\mathcal{M} \times \mathcal{M}$. In this paper we also establish the existence of a large collection of embeddings of $\mathcal{M} \times \mathcal{M}$ and inclusions of $\mathcal{MK}$ in the cubic connectedness locus. The latter restrict to quasiconformal embeddings of every quadratic filled Julia set.

We proceed with a brief overview of our main results. The next section (Section 1) contains a more detailed account of them, including the relevant definitions and statements.

The first objective of this paper is to propose a definition of straightening maps (i.e., to introduce the domains, the target sets, and the maps). We will work in the parameter space $\mathrm{Poly}(d)$ of monic centered polynomials of degree $d \geq 2$ with complex coefficients (i.e., polynomials of the form $z^d + a_{d-2}z^{d-2} + \cdots + a_0$). The *connectedness locus* $C(d)$ *of degree* d is the set formed by all the polynomials $f \in \mathrm{Poly}(d)$ which have connected filled Julia set $K(f)$. We consider a post-critically finite polynomial $f_0 \in C(d)$, with at least one Fatou critical point, and define a straightening map whose domain is, in a certain sense, centered at f_0 and whose target space is introduced via *mapping schemata*. Following Milnor [Mi4], one may associate to f_0 a combinatorial object, called the *reduced mapping schema* T *of* f_0, that encodes the dynamics along the Fatou critical orbits of f_0. *Polynomial maps over* T are certain polynomial dynamical systems which act on a disjoint union of finitely many copies of $\mathbb{C}$. The parameter space formed by all monic centered polynomial maps over T is called *the universal polynomial model space* $\mathrm{Poly}(T)$ *of* T. The notions of Fatou, Julia and filled Julia sets extend to maps in $\mathrm{Poly}(T)$. In $\mathrm{Poly}(T)$, the analogue of the connectedness loci is the *fiberwise connectedness locus* $C(T)$. The fiberwise connectedness locus $C(T)$ will be the target space for a straightening map defined around f_0. The domain $\mathcal{R}(\lambda_0) \subset C(d)$ will be prescribed with the aid of the rational lamination λ_0 of f_0. The definition of the map involves extracting a *polynomial-like map* g *over* T from each polynomial $f \in \mathcal{R}(\lambda_0)$ (i.e., a renormalization procedure). Polynomial-like maps over mapping schemata are a (straightforward) generalization of the classical ones introduced by Douady and Hubbard. The classical notion of hybrid conjugacy between polynomial-like maps also extends easily to our context. Straightening will assign to each polynomial $f \in \mathcal{R}(\lambda_0)$ a map in the fiberwise connectedness locus $C(T)$ which is hybrid equivalent to the polynomial-like map extracted from f. However, in order to avoid ambiguities, we have to introduce "external markings" for polynomial-like maps over T. After carefully introducing markings, we obtain the desired definition of the corresponding straightening map $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \to C(T)$.

Our second objective is to study basic properties of straightening maps. In contrast with the quadratic case, higher degree straightening maps are often discontinuous (see [In4]). Nevertheless, straightening maps are still a natural and useful tool in higher degree polynomial dynamics. Basic questions that one may ask are the following. Are straightening maps injective? What is the image of a given straightening map? When is the domain compact? When is the domain connected? We will show the straightening map $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \to C(T)$, briefly described above, is injective and its image contains all hyperbolic dynamical systems in the fiberwise connectedness locus $C(T)$. Moreover, we give a characterization of straightenings with compact domains. These straightenings include those that arise from "primitive" hyperbolic post-critically finite polynomials.

Straightening maps that arise from "primitive" hyperbolic post-critically finite polynomials are natural candidates to be *onto* the connectedness locus of the corresponding model space. We say that a polynomial $f_0 \in C(d)$ is *primitive* if, for all distinct and bounded Fatou components U, V of f_0, we have that $\overline{U} \cap \overline{V} = \emptyset$. Our third objective is to provide supporting evidence for the following conjecture:

Conjecture. *If* f_0 *is a primitive hyperbolic post-critically finite polynomial with rational lamination* λ_0 *and reduced mapping schema* T*, then* $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \to C(T)$ *is a bijection with compact and connected domain.*

We establish the above conjecture for two large classes of cubic straightening maps: "primitive disjoint type" and "primitive capture type". Moreover, we show that the primitive disjoint type gives rise, via the inverse of straightening, to homeomorphically embedded copies of $\mathcal{M} \times \mathcal{M}$ in the cubic connected locus. Similarly, each primitive capture type, leads to an injective map from the full family of quadratic filled Julia sets $\mathcal{MK}$ into the cubic connectedness locus. The inclusions of $\mathcal{MK}$ in $\mathcal{C}(3)$ have the nice extra property that restrict to a quasiconformal embeddings of every fiber $\{c\} \times K(Q_c)$.

Our results also show that primitive disjoint type straightenings are bijections in any degree (which establishes the existence of embedded copies of $\mathcal{M}^{d-1}$ in $\mathcal{C}(d)$) and, our techniques might generalize to degree $d \geq 4$ versions of primitive capture type straightenings. However, already for cubic polynomials, the above conjecture is open since our techniques do not fully apply for the two remaining types of cubic straightening maps (adjacent and bitransitive).

In [BHe], Buff and Henriksen showed that $K(z^2 + c)$ is quasiconformally embedded in the cubic connectedness locus provided that z^2+c has a non-repelling fixed point. The Buff and Henriksen embeddings are a particular case of the ones obtained via a straightening map of primitive capture type. In [In2], the first author showed that given a degree $d \geq 2$ polynomial $f \in \mathcal{C}(d)$ with filled Julia $K(f)$, then there is a natural embedding of $K(f)$ into the connectedness locus of degree d', for all $d' > d$. Epstein and Yampolsky [EY] obtained embedded copies of $\mathcal{M} \times \mathcal{M}$ by using the intertwining surgery technique.

As mentioned above, one of the main difficulties in the study of higher degree straightening maps stems from the lack of continuity (see [In4]). The other main difficulty stems from the insufficient understanding of the global structure of $\mathcal{C}(d)$, for $d \geq 3$. In fact, the Douady and Hubbard techniques used in the study of the image of quadratic straightening maps break down since they are based on the "topological holomorphy" property of straightening (a very strong form of continuity) and on the knowledge of the global structure of the Mandelbrot set. As suggested by the previous paragraph, the problem of describing the image of higher degree straightening maps has been already addressed in the literature, sometimes in another equivalent language. One approach has been to construct the inverse of some straightening maps via the intertwining construction [EY]. The intertwining technique, which is certainly of intrinsic interest, may only solve the problem for a limited collection of straightening maps. Buff and Henriksen applied holomorphic motions techniques in appropriately chosen one dimensional slices of parameter space. Our approach involves a "combinatorial tuning" technique and obtaining suitable one dimensional restrictions of a given straightening map which behave as in the quadratic case. According to Lyubich [L], quadratic straightening may be also regarded as a holomorphic motion. Nevertheless, we only use the local properties of quadratic straightening to obtain surjectivity rather than the global properties of the holomorphic motions used by Buff and Henriksen.

Acknowledgment. The authors would like to thank Mitsuhiro Shishikura, Peter Haïssinsky and Tomoki Kawahira for valuable discussions.

1. OUTLINE OF RESULTS

The aim of this outline is to provide the reader with a self-contained exposition of the statements of our results. These statements involve several definitions which are introduced in paragraphs 1.1 through 1.9. Paragraphs 1.10 through 1.14 contain the statements of theorems B, C, D, E and F, which are the main results of this paper.

Recall that Poly(d) denotes the space of monic centered polynomials of degree $d \geq 2$. That is, polynomials of the form $z^d + a_{d-2}z^{d-2} + \cdots + a_0$. Therefore, Poly($d$) is naturally identified with $\mathbb{C}^{d-1}$. Also, recall that we denote the connectedness locus by $\mathcal{C}(d)$. According to Branner and Hubbard [BrH], $\mathcal{C}(d)$ is a compact subset of Poly(d). Although the results contained in the body of the paper are stated and proved in further generality, for the sake of simplicity of the exposition, here we restrict our attention to straightening maps that arise from post-critically finite polynomial. Thus:

For rest of this section, we let f_0 be a post-critically finite polynomial with at least one Fatou critical point.

As usual we denote the Julia set by $J(f_0)$, the Fatou set by $F(f_0)$ and the set of critical points by $\mathrm{Crit}(f_0) \subset \mathbb{C}$.

1.1. **Combinatorially renormalizable polynomials.** Given a polynomial $f \in \mathcal{C}(d)$, the *rational lamination λ_f of f* is the equivalence relation in $\mathbb{Q}/\mathbb{Z}$ that identifies two arguments θ, θ' if and only if the external rays of f with arguments θ and θ' land at a common point.

Post-critically finite polynomials in $\mathcal{C}(d)$ are completely determined by their rational laminations [Po]. All the constructions in this paper only depend on a reference rational lamination λ_0. For this outline of results:

We let λ_0 be the rational lamination f_0.

We say that $f \in \mathcal{C}(d)$ is *combinatorially λ_0-renormalizable* if $\lambda_f \supset \lambda_0$. The set formed by all combinatorially λ_0-renormalizable polynomials is denoted by $\mathcal{C}(\lambda_0)$.

1.2. **Mapping schemata.** In [Mi4] a "model" for $\mathcal{C}(\lambda_0)$ is suggested in terms of the reduced mapping schema $T(\lambda_0)$ for f_0.

In general, a *(resp. reduced) mapping schema T* is a triple $(|T|, \sigma, \delta)$ where $|T|$ is a finite set, σ is a map from $|T|$ into itself, and the *degree function δ* is a map $|T|$ into the integers with values (resp. $\delta(v) \geq 2$) $\delta(v) \geq 1$ for all $v \in |T|$.

Most of the mapping schemata considered in this paper are reduced.

The *reduced mapping schema $T(\lambda_0)$ for λ_0 (or f_0)* is $(|T(\lambda_0)|, \sigma_{\lambda_0}, \delta_{\lambda_0})$ where:

(i) $|T(\lambda_0)| = \mathrm{Crit}(f_0) \cap F(f_0)$ is the set of Fatou critical points of f_0.
(ii) $\sigma_{\lambda_0} : |T(\lambda_0)| \to |T(\lambda_0)|$ where $\sigma_{\lambda_0}(v) = {f_0}^{\ell_v}(v)$ if $\ell_v = \min\{k \in \mathbb{N};\ f^k(v) \in |T(\lambda_0)|\}$.
(iii) $\delta_{\lambda_0} : |T(\lambda_0)| \to \mathbb{N}$ where $\delta_{\lambda_0}(v)$ is the local degree of f_0 at v.

The number ℓ_v as above, will be called the *return time of v to $|T(\lambda_0)|$* and will be consistently denoted by ℓ_v.

Note that apparently $T(\lambda_0)$ not only depends on λ_0 but also in f_0. However, we will define $T(\lambda_0)$ purely in terms of λ_0 in Section 3.3.

1.3. **Universal polynomial model space.** Each reduced mapping schema $T = (|T|, \sigma, \delta)$ determines a complex affine space $\mathrm{Poly}(T)$ called the *universal polynomial model space for T*. More precisely, $\mathrm{Poly}(T)$ consists of all maps f from $|T| \times \mathbb{C}$ to itself such that the restriction of f to each component $\{v\} \times \mathbb{C}$ is a monic centered polynomial map f_v of degree $\delta(v)$, taking values in $\{\sigma(v)\} \times \mathbb{C}$. For short we say that f is a *polynomial map over T*. Given $f \in \mathrm{Poly}(T)$, the *filled Julia set $K(f)$* of f is the set of points in $|T| \times \mathbb{C}$ with precompact forward orbit. The boundary $\partial K(f)$ is called the *Julia set $J(f)$ of f*. We say that $K(f)$ is *fiberwise connected* if the intersection of $K(f)$ with every fiber $\{v\} \times \mathbb{C}$ is connected. The subset of $\mathrm{Poly}(T)$ formed by the maps f with fiberwise connected filled Julia set $K(f)$ is called the *connectedness locus $\mathcal{C}(T)$ for T*. It follows that $\mathcal{C}(T)$ is a compact subset of $\mathrm{Poly}(T)$. We will denote the v-fiber of the filled Julia set $K(f)$ by $K(f, v)$. Also, we will abuse of notation and sometimes regard $f \in \mathrm{Poly}(T)$ as a collection $f = (f_v : \mathbb{C} \to \mathbb{C})_{v \in |T|}$ of polynomials.

1.4. **Polynomial-like maps over schemata.** Now we generalize the notion of polynomial-like maps introduced by Douady and Hubbard [DH2].

Consider a mapping schema $T = (|T|, \sigma, \delta)$. Let $U' \Subset U$ be open subsets contained in $|T| \times \mathbb{C}$ which are fiberwise topological disks. That is, for all $v \in |T|$ we have that $U' \cap (\{v\} \times \mathbb{C}) = \{v\} \times U'_v$ and $U \cap (\{v\} \times \mathbb{C}) = \{v\} \times U_v$ for some topological disks $U'_v, U_v \subset \mathbb{C}$. A proper and holomorphic skew product over σ,

$$\begin{array}{cccc} g: & U' & \to & U \\ & (v,z) & \to & (\sigma(v), g_v(z)) \end{array}$$

is called a *polynomial-like map over* T if the degree of $g_v : U'_v \to U_{\sigma(v)}$ is $\delta(v)$, for every $v \in |T|$. The *filled Julia set* $K(g)$ is the set of all $(v,z) \in U'$ such that $g^n((v,z))$ is well defined, for all $n \in \mathbb{N}$. Again we will abuse of notation and sometimes simply identify g with the collection $(g_v : U'_v \to U_{\sigma(v)})_{v \in |T|}$. The v-fiber of $K(g)$ will be denoted by $K(g,v)$.

Simple examples of polynomial-like maps over a reduced mapping schema T are obtained from polynomial maps in $\mathcal{C}(T)$ after restriction to an appropriate neighborhood of their filled Julia sets.

Two polynomial-like maps g_0 and g_1 over a mapping schema T with are said to be *hybrid equivalent* if there exists a fiberwise quasiconformal map ψ defined on a neighborhood of $K(g_0)$ such that $\psi \circ g_0 = g_1 \circ \psi$ and

$$\frac{\partial \psi}{\partial \bar{z}}(v,z) \equiv 0 \text{ a.e. on } K(g_0).$$

1.5. **External markings.** We will show that every polynomial-like map g over a reduced mapping schema T with fiberwise connected Julia set is hybrid equivalent to a polynomial map over T. However, in order to avoid ambiguities we are forced either to work with the moduli space of polynomial maps over T (affine conjugacy classes of such maps) or to introduce *external markings*. For our purpose, the latter will be more convenient. Roughly speaking, an external marking of such a polynomial-like map g over T is a collection of accesses, one per fiber, to an appropriate periodic or preperiodic point in $J(g)$.

Definition 1.1. Let $g : U' \to U$ be a polynomial-like map over a reduced mapping schema $T = (|T|, \sigma, \delta)$ with connected filled Julia set. A *path to* $K(g)$ is a path $\gamma : [0,1] \to U'$ such that $\gamma((0,1]) \subset U' \setminus K(g)$ and $\gamma(0) \in J(g)$. We say two paths γ_0 and γ_1 to $K(f)$ are *homotopic* if there exists a continuous map $\tilde{\gamma} : [0,1] \times [0,1] \to U'$ such that

- $t \mapsto \tilde{\gamma}(s,t)$ is a path to $K(g)$ for any $s \in [0,1]$;
- $\tilde{\gamma}(0,t) = \gamma_0(t)$ and $\tilde{\gamma}(1,t) = \gamma_1(t)$;
- $\tilde{\gamma}(s,0) = \gamma_0(0)$ for any $s \in [0,1]$.

An *access to* $K(g)$ is a homotopy class of paths to $K(g)$.

Definition 1.2. Let $g : U' \to U$ be a polynomial-like map over a reduced mapping schema $T = (|T|, \sigma, \delta)$ with $K(g)$ connected. An *external marking of* g is a collection $\Gamma = (\Gamma_v)_{v \in |T|}$ where each Γ_v is an access in $\{v\} \times \mathbb{C}$ and such that Γ is forward invariant in the following sense. For every $v \in |T|$ and every representative $\gamma_v \subset (\{v\} \times \mathbb{C}) \cap U'$ of Γ_v, the connected component of $g(\gamma_v) \cap U'$ that contains a point of $J(g)$ is a representative of $\Gamma_{\sigma(v)}$.

An *externally marked polynomial-like map over* T is a pair (g, Γ) of a polynomial-like map over T and an external marking of it.

1.6. **Standard external marking of polynomial maps in** $\mathrm{Poly}(T)$**.** Polynomial maps over a reduced mapping schema T are naturally endowed with a standard marking which we introduce with the aid of appropriately defined Böttcher coordinates. More precisely, given

$f \in \mathrm{Poly}(T)$, the standard arguments for polynomials (e.g., see [Mi3]) easily generalize to prove that there exists a *Böttcher coordinate* φ (at infinity) for f. That is, there exists a neighborhood U of $|T| \times \{\infty\}$ in $|T| \times \hat{\mathbb{C}}$ and a conformal map $\varphi : U \to |T| \times \hat{\mathbb{C}}$ of the form $\varphi(v, z) = (v, \varphi_v(z))$ such that $\varphi \circ f(v, z) = (\sigma(v), (\varphi_v(z))^{\delta(v)})$ and φ is asymptotic to the identity as $z \to \infty$ on each fiber.

Let $\Delta = \{z \in \mathbb{C}; |z| < 1\}$. If $f \in \mathcal{C}(T)$, then a Böttcher coordinate can be extended uniquely to a conformal isomorphism

$$\varphi : (|T| \times \mathbb{C}) \setminus K(f) \to |T| \times \left(\mathbb{C} \setminus \overline{\Delta}\right),$$

which we also denote by φ. In this case, φ is uniquely determined by $f \in \mathcal{C}(T)$.

For $(v, \theta) \in |T| \times \mathbb{R}/\mathbb{Z}$, define the *external ray for* f by

$$R_f(v, \theta) = \{\varphi^{-1}(v, r \exp(2\pi i\theta));\ 1 < r < \infty\} \quad (v \in |T|,\ \theta \in \mathbb{R}/\mathbb{Z}).$$

We say that $R_f(v, \theta)$ *lands at* $(v, z) \in J(f)$ if

$$\lim_{r \searrow 1} \varphi^{-1}(v, r \exp(2\pi i\theta)) = (v, z).$$

The landing theorems easily generalize to this context (e.g., see [Mi3]). In particular, external rays with arguments in $\mathbb{Q}/\mathbb{Z}$ always land (at eventually periodic points).

For $f \in \mathcal{C}(T)$, the collection

$$(R_f(v, 0))_{v \in |T|}$$

of the external rays of angle 0 naturally induces an external marking of any polynomial-like restriction of f. We call it the *standard external marking of* f. Figure 1 is an example of external markings of polynomial of degree 4. It has three external markings, each corresponds to some invariant external ray. The standard marking corresponds to the 0-ray (i.e., the horizontal one).

1.7. **Straightening of polynomial-like maps over reduced schemata.** Consider two externally marked polynomial-like maps $(g_0 : U_0' \to U_0, \Gamma_0)$, $(g_1 : U_1' \to U_1, \Gamma_1)$ over T with connected filled Julia sets and assume there exists a topological conjugacy $\psi : U_0 \to U_1$ between g_0 and g_1. We say ψ *respects external markings* if $\Gamma_1 = ([\psi \circ \gamma_v])_{v \in |T|}$ where $(\gamma_v)_{v \in |T|}$ is a representative of Γ_0. Note that this definition is independent of the choice of representative.

Section 2 contains the following generalization of Douady and Hubbard straightening theorem to our context.

Theorem A (Straightening). *Let T be a reduced mapping schema and consider a polynomial-like map g over T. Then, there exists a polynomial map $f \in \mathrm{Poly}(T)$ hybrid conjugate to g. Moreover,*

(i) *If $K(g)$ is fiberwise connected, then $f \in \mathcal{C}(T)$ and f is unique up to affine conjugacy.*

(ii) *If $K(g)$ is fiberwise connected and g is externally marked by Γ, then for exactly one $f \in \mathcal{C}(T)$ there exists a hybrid conjugacy that respects the external markings Γ and Γ', where Γ' is the standard external marking.*

Based on the previous result which "straightens" a single polynomial-like map, we will define "straightening maps" from an appropriately defined domains contained in the connectedness locus.



FIGURE 1. The invariant rays for a quartic polynomial $z^4 - 1$. Different invariant rays represent different external markings.

1.8. **Renormalizable polynomials.** Recall from Section 1.1 that $C(\lambda_0)$ denotes the λ_0-combinatorially renormalizable locus. Under certain conditions it will be possible to extract a polynomial-like map over $T(\lambda_0)$ from a given $f \in C(\lambda_0)$. In order to be more precise, below, for every $v \in |T(\lambda_0)|$, we identify a subset $K_v(f)$ of $K(f)$ which we call the *v-small filled Julia set*.

Given $\theta, \theta' \in \mathbb{Q}/\mathbb{Z}$ so that the corresponding rays $R_{f_0}(\theta), R_{f_0}(\theta')$ land at a common point (i.e., θ, θ' are λ_0-equivalent) let $\text{Sector}_{f_0}(\theta, \theta'; v)$ be the connected component of $\mathbb{C} \setminus \overline{(R_{f_0}(\theta) \cup R_{f_0}(\theta'))}$ that contains v. If $f \in C(\lambda_0)$, then we denote by $\text{Sector}_f(\theta, \theta'; v)$ the connected component of $\mathbb{C} \setminus \overline{(R_f(\theta) \cup R_f(\theta'))}$ such that, for all $t \in \mathbb{R}/\mathbb{Z}$,

$$R_{f_0}(t) \subset \text{Sector}_{f_0}(\theta, \theta'; v) \text{ if and only if } R_f(t) \subset \text{Sector}_f(\theta, \theta'; v).$$

Now we define,

$$K_v(f) = K(f) \cap \bigcap_{\theta \sim_{\lambda_0} \theta'} \overline{\text{Sector}_f(\theta, \theta'; v)}.$$

It follows that $K_v(f)$ is connected and $f^{\ell_v}(K_v(f)) = K_{\sigma_{\lambda_0}(v)}(f)$ (see Proposition 3.7).

We say that $f \in C(\lambda_0)$ is λ_0-*renormalizable* if for all $v \in \text{Crit}(f_0) \cap F(f_0)$ there exist topological disks $U'_v \Subset U_v$ such that $K_v(f) \subset U'_v$ and $f^{\ell_v} : U'_v \to U_{\sigma_{\lambda_0}(v)}$ is a proper map of degree $\delta_{\lambda_0}(v)$. We denote the set of λ_0-renormalizable polynomials by $\mathcal{R}(\lambda_0)$.

Consider $f \in \mathcal{R}(\lambda_0)$ and, using the notation above, we let

$$U' = \{(v, z);\ z \in U'_v\}$$

$$U = \{(v, z);\ z \in U_v\}$$

and $g_v = f^{\ell_v}$. It follows that

$$\begin{array}{rccl} g: & U' & \to & U \\ & (v, z) & \mapsto & (\sigma_{\lambda_0}(v), g_v(z)) \end{array}$$

is a polynomial-like map over $T(\lambda_0)$ with (fiberwise connected) filled Julia set

$$K(g) = \{(v, z);\ z \in K_v(f)\}.$$

We say that g is a λ_0-*renormalization* of f. Note that g is uniquely defined over $K(g)$, however there is a choice involved for the domain U'.

1.9. **Internal angle systems and induced external markings.** We consistently mark the polynomial-like maps over $T(\lambda_0)$ extracted from maps in $\mathcal{R}(\lambda_0)$. More precisely, we introduce *internal angle systems* for f_0 and describe how an internal angle system determines an external marking for a λ_0-renormalization of every $f \in \mathcal{R}(\lambda_0)$.

For every $v \in |T(\lambda_0)|$ denote by γ_v the boundary of the Fatou component of f_0 that contains v. Since f_0 is post-critically finite, γ_v is a Jordan curve (e.g., see [Mi3]). Moreover, there exists a collection $\alpha = (\alpha_v : \gamma_v \to \mathbb{R}/\mathbb{Z})_{v \in |T(\lambda_0)|}$ of homeomorphisms such that:

$$\alpha_{\sigma_{\lambda_0}(v)}(f_0{}^{\ell_v}(z)) = \delta_{\lambda_0}(v)\alpha_v(z)$$

for all $z \in \gamma_v$.

We call $\alpha = (\alpha_v)_{v \in |T(\lambda_0)|}$ an *internal angle system* for f_0.

An internal angle system determines an external marking of any λ_0-renormalization of every $f \in \mathcal{R}(\lambda_0)$. For each $v \in |T(\lambda_0)|$ choose an argument θ_v so that the external ray of f_0 with argument θ_v lands at $\alpha_v^{-1}(0)$. Given a λ_0-renormalization g of a polynomial $f \in \mathcal{R}(\lambda_0)$, let Γ_v be the access with representative the connected component of $\overline{R_f(\theta_v)} \cap U'_v$ that contains a point of $K_v(f)$. We say that $\Gamma = (\Gamma_v)_{v \in |T(\lambda_0)|}$ is the *external marking of g determined by the internal angle system* $(\alpha_v : \gamma_v \to \mathbb{R}/\mathbb{Z})_{v \in |T(\lambda_0)|}$. This external marking is independent of the choices involved (see Section 3.3).

1.10. **Straightening map and injectivity.** Finally we are ready to define the straightening maps under consideration. More precisely, as a consequence of Theorem A we obtain the following result.

Corollary. *Given an internal angle system α for f_0 and $f \in \mathcal{R}(\lambda_0)$, there exists a unique map $P \in C(T(\lambda_0))$ such that there exists a hybrid equivalence between P and a λ_0-renormalization g of f, respecting external markings, where the external marking of g is the one determined by α and the external marking of P is the standard external marking.*

With the notation of the above corollary, given an internal angle system α for f_0, we say that the *associated straightening map* $\chi_{\lambda_0} : C(\lambda_0) \to C(T(\lambda_0))$ is the function defined by $\chi_{\lambda_0}(f) = P$. For example, consider $f_0(z) = z^4 - 1$ as in Figure 2. There are three polynomials f_1, f_2, and $f_3 \in \mathcal{R}(\lambda_0)$ such that $\chi_{\lambda_0}(f_i)$ is affinely conjugate to f_0 depending on the choice of the external marking of f_0. Figure 3 shows two of f_i's and the other one is just the complex conjugate of the second one.

In Section 5 we prove the following result.


(a) The Julia set and Yoccoz puzzles (b) The rational lamination

FIGURE 2. The Julia set and Yoccoz puzzles, and the rational lamination for $z^4 - 1$.

Theorem B (Injectivity of Straightening)**.** *Consider an internally angled post-critically finite polynomial* f_0. *Let* $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \to C(T(\lambda_0))$ *be the associated straightening map. Then,* χ_{λ_0} *is injective.*

1.11. **Onto hyperbolic maps.** To study the basic properties of the image of general straightening maps, we employ a "combinatorial tuning procedure" which is discussed in Section 6.

Theorem C. *Let* f_0 *be an internally angled post-critically finite polynomial such that* $\mathcal{R}(\lambda_0) \neq \emptyset$ *and* $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \to C(T(\lambda_0))$ *be the associated straightening map. Denote by* $\operatorname{Hyp} C(T(\lambda_0))$ *the set of hyperbolic maps contained in* $C(T(\lambda_0))$. *Then* $\chi_{\lambda_0}(\mathcal{R}(\lambda_0)) \supset \operatorname{Hyp} C(T(\lambda_0))$ *and*

$$\chi_{\lambda_0} : \chi_{\lambda_0}^{-1}(\operatorname{Hyp} C(T(\lambda_0)) \to \operatorname{Hyp} C(T(\lambda_0))$$

is biholomorphic.

An equivalent condition for $\mathcal{R}(\lambda_0) \neq \emptyset$ is given in Proposition 6.7 (see also Theorem D).

1.12. **Compactness.** Compactness properties of the domain of a straightening map are useful to further study its image. Recall that we say that a post-critically finite polynomial $f \in C(d)$ is *primitive* if f has at least one periodic critical point and, for every pair of distinct bounded Fatou components V_1, V_2 of f_0, we have that $\partial V_1 \cap \partial V_2 = \emptyset$.

Theorem D. *Let* f_0 *be a post-critically finite polynomial. Then the following statements are equivalent:*

(i) f_0 *is primitive.*
(ii) $C(\lambda_0) = \mathcal{R}(\lambda_0) \neq \emptyset$.
(iii) $\mathcal{R}(\lambda_0)$ *is compact and non-empty.*



FIGURE 3. Tunings of $z^4 - 1$ with itself with different external markings, given by the external rays of angle 0 and $1/3$.

1.13. **Cubic reduced mapping schemata.** There are exactly four types of cubic reduced mapping schemata.

- $T_{\mathrm{adj}} = (\{0\}, \sigma, \delta)$, where $\sigma(0) = 0$ and $\delta(0) = 3$, is called a reduced mapping schema of *adjacent type*. For example, if f_0 is polynomial of the form $z^3 + c$ for which the critical point $z = 0$ is periodic, then the reduced mapping schema $T(\lambda_0)$ is T_{adj}. Observe that $\mathrm{Poly}(T_{\mathrm{adj}}) = \mathrm{Poly}(3)$ and $\mathcal{C}(T_{\mathrm{adj}}) = \mathcal{C}(3)$.
- $T_{\mathrm{bit}} = (\{v_0, v_1\}, \sigma, \delta)$, where $\sigma(v_j) = v_{1-j}$ for $j = 0, 1$, and δ is constant $(= 2)$, is called a reduced mapping schema of *bitransitive type*. If f_0 is a cubic polynomial with distinct critical points v_0, v_1 lying in the same periodic orbit, then $T(\lambda_0) = T_{\mathrm{bit}}$. Note that $\mathrm{Poly}(T_{\mathrm{bit}})$ can be identified with the family of biquadratic polynomials $\mathrm{Poly}(2 \times 2) = \{(z^2 + c_0)^2 + c_1;\ (c_0, c_1) \in \mathbb{C}^2\} \subset \mathrm{Poly}(4)$. Thus the connectedness locus of $\mathrm{Poly}(T_{\mathrm{bit}})$ is identified with $\{(c_0, c_1) \in \mathbb{C}^2;\ (z^2 + c_0)^2 + c_1 \in \mathcal{C}(4)\}$.
- $T_{\mathrm{cap}} = (\{v_0, v_1\}, \sigma, \delta)$, where $\sigma(v_j) = v_0$ for $j = 0, 1$, and δ is constant $(= 2)$, is called a reduced mapping schema of *capture type*. If f_0 is a cubic polynomial with distinct

critical points v_0, v_1 such that v_0 is periodic and v_1 is not periodic but eventually lands in the orbit of v_0, then $T(\lambda_0) = T_{\mathrm{cap}}$. In this case, $\mathrm{Poly}(T)$ is naturally identified with $\mathbb{C}^2$. In fact, given a map $f : \{v_0, v_1\} \times \mathbb{C} \to \{v_0, v_1\} \times \mathbb{C}$ in $\mathrm{Poly}(T)$ there exists (c_0, c_1) such that $f(v_j, z) = (v_0, z^2 + c_j)$. Note that $K(f)$ is fiberwise connected if and only if $c_0 \in \mathcal{M}$ and $c_1 \in K(z^2 + c_0)$. Therefore,

$$\mathcal{C}(T_{\mathrm{cap}}) = \mathcal{MK} = \{(c, z) \in \mathbb{C}^2;\ c \in \mathcal{M}, z \in K(z^2 + c)\}.$$

- $T_{\mathrm{dis}} = (\{v_0, v_1\}, \sigma, \delta)$ where $\sigma(v_j) = v_j$ for $j = 0, 1$, and δ is constant $(= 2)$, is called a reduced mapping schema of *disjoint type*. If f_0 is a cubic polynomial with distinct critical points v_0, v_1 such that both critical points are periodic but belong to different orbits, then $T(\lambda_0) = T_{\mathrm{dis}}$. It follows that $\mathrm{Poly}(T_{\mathrm{dis}})$ is identified with $\mathrm{Poly}(2) \times \mathrm{Poly}(2)$ and $\mathcal{C}(T_{\mathrm{dis}}) = \mathcal{M} \times \mathcal{M}$.



T_{adj}: adjacent type



T_{bit}: bitransitive type



T_{cap}: capture type



T_{dis}: disjoint type

1.14. **Capture and Disjoint type straightenings.** Using the compactness given by Theorem D we are able to extend Theorem C and describe the image of two classes straightening maps centered at a primitive hyperbolic post-critically finite polynomial f_0. The first class is the generalization of the notion of *disjoint type*; described above in the context of cubic polynomials. Namely, the reduced mapping schema T consists of $d - 1$ elements which are pointwise fixed by the schema map. Equivalently, f_0 has exactly $d - 1$ superattracting periodic orbits. Note that the universal polynomial model space for T, consists of ordered $(d - 1)$-tuples of monic centered quadratic polynomials which act on $d - 1$ copies of $\mathbb{C}$. Thus, the corresponding connectedness locus is $\mathcal{M}^{d-1}$.

Theorem E. *Let $f_0 \in \mathcal{C}(d)$ be a polynomial with exactly $d - 1$ superattracting periodic orbits which is primitive. Then, given any internal angle system, the associated straightening map*

$$\begin{array}{rccc} \chi: & \mathcal{R}(\lambda_{f_0}) & \to & \mathcal{M}^{d-1} \\ & f & \mapsto & (\chi_1(f), \ldots, \chi_{d-1}(f)), \end{array}$$

is a homeomorphism. Moreover, for all $1 \leq i_1 < \cdots < i_k \leq d - 1$ and all $c_1, \ldots, c_k \in \mathcal{M}$, the set

$$\{f \in \mathcal{R}(\lambda_0);\ \chi_{i_j}(f) = c_j \text{ for all } j = 1, \ldots, k\}$$

is contained in a codimension k complex submanifold $\mathcal{S}$ of $\mathrm{Poly}(d)$. *Furthermore, if $k = d - 2$, then $\chi_j : \mathcal{S} \cap \mathcal{R}(\lambda_0) \to \mathcal{M}$ extends to a quasiconformal map in a neighborhood of $\mathcal{S} \cap \mathcal{R}(\lambda_0)$.*

The second class is that of cubic straightening of capture type.

Theorem F. *Let $f_0 \in \mathcal{C}(3)$ be an internally angled primitive hyperbolic post-critically finite polynomial with rational lamination λ_0 and reduced mapping schema of capture type. Then $\mathcal{R}(\lambda_0)$ is connected, and the associated straightening map*

$$\begin{array}{rccc} \chi: & \mathcal{R}(\lambda_0) & \to & \mathcal{MK} \\ & f & \mapsto & (\chi_1(f), \chi_2(f)), \end{array}$$

is a bijection. Moreover, given $c \in \mathcal{M}$, the set

$$\{f \in \mathcal{R}(\lambda_0);\ \chi_1(f) = c\}$$

is contained in a one dimensional complex analytic space $\mathcal{S}_c$ which is locally irreducible. Furthermore,

$$\chi_2 : \mathcal{R}(\lambda_0) \cap \mathcal{S}_c \to K(z^2 + c)$$

extends to a quasiconformal map from a neighborhood of $\mathcal{R}(\lambda_0) \cap \mathcal{S}_c$ onto a neighborhood of $K(z^2 + c)$.

Note that, in the previous theorem, $\mathcal{S}_c$ might be a singular analytic space. In order have well defined quasiconformal maps with domain in $\mathcal{S}_c$ we need to specify a conformal structure. In fact, there is only one such structure compatible with the embedding, since around each singular point of $\mathcal{S}_c$ there exists a punctured neighborhood U (in $\mathcal{S}_c$) and a conformal isomorphism from the punctured disk onto U which has a unique continuous extension.

2. Polynomial-like maps

The aim of this section is to prove Theorem A (see [DH1]). That is, to generalize Douady and Hubbard straightening Theorem to the context of polynomial-like maps over reduced mapping schemata.

2.1. **Proof of the straightening theorem.**

Let g be a polynomial-like map over a reduced mapping schema T. Following the ideas of Douady and Hubbard, we "paste" g with the dynamics of the polynomial map over T given by $f_0(v, z) = (\sigma(v), z^{\delta(v)})$.

Let $\overline{\Delta}(r) = \{|z| \le r\}$ denote the closed disk of radius $r > 0$. Choose $R' > R > 1$. Let $V = |T| \times (\hat{\mathbb{C}} \setminus \overline{\Delta}(R))$ and

$$V' = f_0(V) \cup \Big((|T| \setminus \sigma(|T|)) \times (\hat{\mathbb{C}} \setminus \overline{\Delta}(R'))\Big).$$

Restricting g to a smaller domain, if necessary, we may assume the domain U' and the range U have C^1 boundaries. Take a C^1 diffeomorphism h between $A = U \setminus U'$ and $B = V \setminus V'$ such that $h(v, z) = (v, h_v(z))$ for some h_v and $h(g(v, z)) = f_0(h(v, z))$ for all $z \in \partial U'$. Then consider the C^1 manifold $S = (U \sqcup V)/h$ and the map $\tilde{f} : S \to S$ defined by

$$\tilde{f}(v, z) = \begin{cases} g(v, z) & \text{if } (v, z) \in U', \\ f_0(v, z) & \text{if } (v, z) \in V. \end{cases}$$

Applying the standard pull-back argument, we obtain a $\tilde{f}$-invariant almost complex structure μ_0 on S. From the measurable Riemann mapping theorem, we obtain a biholomorphic map from (S, μ_0) onto $|T| \times \hat{\mathbb{C}}$ which conjugates $\tilde{f}$ with a polynomial map f over $|T|$. Moreover, since μ_0 was obtained via the pull-back argument, it follows that g and f are hybrid conjugate.

The uniqueness part of Theorem A follows from the proposition below.

Proposition 2.1. *If two polynomial maps $f_0, f_1 \in \mathrm{Poly}(T)$ over a reduced mapping schema T are hybrid equivalent, then they are affine conjugate.*

Moreover, if $f_0, f_1 \in C(T)$ and a hybrid conjugacy respects the standard external markings, then $f_0 = f_1$ and the affine conjugacy is the identity.

Before proving the proposition let us give a short discussion about external markings and affine conjugacies. Below, we show that, modulo affine conjugacy, there is only one external marking.

Given a polynomial map $f \in \mathcal{C}(T)$ consider an external marking Γ for (a polynomial-like restriction of) f. Since every access contains a unique external ray, by Lindelöf's theorem, we can choose a collection of external rays $(R_f(v, \theta_v))_{v \in |T|}$ as a representative of Γ. It follows that,

$$\delta(v)\theta_v = \theta_{\sigma(v)}. \tag{1}$$

Observe that there are only finitely many collections of angles (θ_v) satisfying (1).

Now let

$$A(v, z) = (v, e^{2\pi i \theta_v} z),$$

and $\hat{f} = A^{-1} \circ f \circ A$. It is easy to check that $\hat{f} \in \mathcal{C}(T)$ and $A(R_{\hat{f}}(v, \theta)) = R_f(v, \theta + \theta_v)$. In particular, A maps the standard external marking for $\hat{f}$ to the external marking $(R_f(v, \theta_v))$.

On the other hand, it is easy to check that if a collection $(\theta_v)_{v \in |T|}$ of angles satisfies (1), then $(R_f(v, \theta_v))$ defines an external marking Γ for f.

Now we prove the proposition and, therefore, we finish the proof of Theorem A.

Proof. As in the statement of the proposition we consider two polynomial maps f_0, f_1 over T with hybrid conjugate polynomial-like restrictions $f_j : U'_j \to U_j$ where $j = 0, 1$. Let us denote the hybrid conjugacy $\psi : U_0 \to U_1$. Replacing f_1 by an affinely conjugate polynomial map we may assume that ψ respects the standard external markings.

Define a homeomorphism Φ by

$$\Phi(v, z) = \begin{cases} \psi(v, z) & \text{if } (v, z) \in K(f_0), \\ \varphi_{f_1}^{-1} \circ \varphi_{f_0}(v, z) & \text{otherwise.} \end{cases}$$

Clearly, $\Phi \circ f_0 = f_1 \circ \Phi$.

As in the proof of [DH2, Proposition 6], since ψ respects the standard external markings, it follows that Φ is biholomorphic, hence affine. The affine map Φ conjugates two monic centered polynomial maps over T and it is asymptotic to the identity at infinity in every fiber. Therefore, Φ must be the identity. □

3. LAMINATIONS

3.1. **Laminations.** Laminations were introduced to the context of polynomial dynamics in the early 1980's by Thurston [Th]. Since then they have been proven to be an useful object to encode the landing pattern of external rays. In this section, we briefly review the basic notions and results about laminations. For our purposes we will need to consider the well known d-invariant laminations as well as forward invariant laminations with finite support. The latter type of laminations are used to define "combinatorial Yoccoz puzzles" (compare with [In3]).

For an integer $d > 0$, we let

$$\begin{array}{rccc} m_d : & \mathbb{R}/\mathbb{Z} & \to & \mathbb{R}/\mathbb{Z} \\ & \theta & \mapsto & d\theta. \end{array}$$

Two subsets $A, B \subset E$ are said to be *unlinked* if B is contained in a component of $\mathbb{R}/\mathbb{Z} \setminus A$ (it is equivalent to A being contained in a component of $\mathbb{R}/\mathbb{Z} \setminus B$). Note that A, B are unlinked if and only if the Euclidean (or hyperbolic) convex hulls in $\overline{\Delta} = \{|z| \leq 1\}$ of $\exp(2\pi i A)$ and $\exp(2\pi i B)$ are disjoint.

Definition 3.1 (Laminations without dynamics)**.** Let $E \subset \mathbb{R}/\mathbb{Z}$. An equivalence relation λ on E is called a *lamination on E* if the following three conditions hold.

(i) λ is closed in $E \times E$.
(ii) Every equivalence class is finite.
(iii) Equivalence classes are pairwise unlinked.

The *support* $\mathrm{supp}(\lambda)$ of λ is the union of all non-trivial λ-equivalence classes.

A lamination on $\mathbb{Q}/\mathbb{Z}$ (resp. $\mathbb{R}/\mathbb{Z}$) is called a *rational* (resp. *real*) *lamination*.

We will be mainly concerned with laminations that are compatible with the action of m_d.

Definition 3.2 (Forward invariant and d-invariant laminations)**.** Let $d > 1$ be an integer and $E \subset \mathbb{R}/\mathbb{Z}$. Let λ and λ' be laminations on E and $E' = m_d^{-1}(E)$, respectively.

We say that $(m_d)_*\lambda = \lambda'$ if for any λ-equivalence class A,

(i) $m_d(A)$ is a λ'-equivalence class.
(ii) $m_d : A \to m_d(A)$ is *consecutive preserving*, i.e., for any component (θ, θ') of $\mathbb{R}/\mathbb{Z} \setminus A$, $(d\theta, d\theta')$ is a component of $\mathbb{R}/\mathbb{Z} \setminus m_d(A)$.

A lamination λ is said to be *forward invariant* if $(m_d)_*\lambda = \lambda$. A forward invariant lamination λ on a m_d-completely invariant set $E \subset \mathbb{R}/\mathbb{Z}$ is called *d-invariant* or simply *invariant*.

3.2. **Invariant rational laminations: unlinked classes, fibers and critical elements.**

Invariant rational laminations will be particularly useful in this paper. Recall that $\mathcal{C}(d)$ denotes connectedness locus in the space of monic centered degree $d \geq 2$ polynomials $\mathrm{Poly}(d)$. Given a monic centered polynomial $f \in \mathcal{C}(d)$ the *rational lamination λ_f of f* is the equivalence relation on $\mathbb{Q}/\mathbb{Z}$ which identifies two rational arguments if and only if the corresponding rays land at a common point. According to [Ki], an equivalence relation λ on $\mathbb{Q}/\mathbb{Z}$ is the rational lamination of some $f \in \mathcal{C}(d)$ if and only if λ is a d-invariant rational lamination.

Rational laminations allow us to establish when distinct polynomial share certain dynamical features. Therefore, it is natural to define some subsets of dynamical and parameter space in terms of rational laminations. Given a d-invariant rational lamination λ, let

$$\mathcal{C}(\lambda) = \{f \in \mathcal{C}(d); \lambda_f \supset \lambda\}.$$

Note that this set is always non-empty [Ki].

The sets $\mathcal{C}(\lambda)$ have already deserved a lot of attention. When λ is the rational lamination of a polynomial with all cycles repelling (equivalently, λ is maximal with respect to the partial order on d-invariant rational laminations), $\mathcal{C}(\lambda)$ is a "combinatorial class". In degree 2, the MLC conjecture asserts that combinatorial classes are singletons. In degree 3, there are some combinatorial classes which are non-trivial continua [He]. However, it is natural to conjecture that in any degree, each combinatorial classes is contained in an analytic subspace of $\mathrm{Poly}(d)$ of codimension (at least) 1 (see [In2]).

This paper is mainly concerned with $\mathcal{C}(\lambda)$ for invariant rational laminations λ which are not maximal. In degree 2, following Douady and Hubbard, these sets coincide with homeomorphic copies of the Mandelbrot contained contained in the Mandelbrot set $\mathcal{M}$. In order to compare $\mathcal{C}(\lambda)$ with its model we employ a renormalization procedure. A first step towards renormalization is to cut dynamical space into sectors and the filled Julia set into pieces.

Definition 3.3 (Admissible)**.** Consider a polynomial $f \in \mathcal{C}(d)$. We say that a lamination λ on $E \subset \mathbb{Q}/\mathbb{Z}$ is *admissible for f* if for every pair of distinct arguments θ and θ' which are λ-related, we have that they are also λ_f-related.

Note that $\mathcal{C}(\lambda)$ exactly consists of all the polynomials for which the invariant rational lamination λ is admissible. Admissible laminations will allow us to cut the dynamical space into sectors and the filled Julia set into fibers (compare with [Sch]).

Definition 3.4 (Sectors and fibers)**.** Let $f \in \mathcal{C}(d)$ and λ be a lamination on $E \subset \mathbb{Q}/\mathbb{Z}$ that is admissible for f. Consider a set $L \subset \mathbb{R}/\mathbb{Z}$ which is unlinked with every non-trivial λ-class. If θ and θ' are distinct λ-equivalent arguments, we denote by

$$\operatorname{Sector}(\theta, \theta'; L)$$

the connected component of $\mathbb{C} \setminus \overline{(R_f(\theta) \cup R_f(\theta'))}$ which contains every external ray $R_f(t)$ with $t \in L$. Moreover, we define the *λ-fiber* of L by:

$$K_f(L) = K(f) \cap \bigcap_{\theta \sim_\lambda \theta', \theta \neq \theta'} \overline{\operatorname{Sector}(\theta, \theta'; L)}.$$

Our main interest now is when λ is an invariant rational lamination and the sets L above are, in a certain sense, as large as possible. That is, when L are "unlinked classes".

Definition 3.5 (Unlinked classes)**.** Let λ be a d-invariant rational lamination. We say $\theta, \theta' \in \mathbb{R}/\mathbb{Z} \setminus \mathbb{Q}/\mathbb{Z}$ are λ-unlinked if $\theta = \theta'$ or for any λ-equivalence class A, θ and θ' lie in the same component of $\mathbb{R}/\mathbb{Z} \setminus A$.

Observe that, by definition, unlinked classes are contained in $\mathbb{R}/\mathbb{Z} \setminus \mathbb{Q}/\mathbb{Z}$. Moreover, the λ-unlinked relation is in fact an equivalence relation. Infinite λ-unlinked classes are closely related to the concept of gaps introduced by Thurston.

In order to visualize the unlinked classes of an invariant rational lamination it is convenient to consider the real extension of a rational lamination.

Definition 3.6 (Real extension)**.** Given a lamination λ on $E \subset \mathbb{R}/\mathbb{Z}$ the *real extension $\hat{\lambda}$ of λ* is the smallest equivalence relation that contains the closure $\overline{\lambda}$ of λ in $\mathbb{R}/\mathbb{Z} \times \mathbb{R}/\mathbb{Z}$.

According to [Ki], the real extension of an invariant rational lamination is an invariant real lamination.

For example, consider a polynomial f with locally connected Julia set and such that the following technical condition is satisfied. No critical point with infinite forward orbit lies in the boundary of a bounded Fatou component (e.g., f is post-critically finite). Then, the real extension of the rational lamination λ_f encodes the landing pattern of *all* the external rays of f. That is $\widehat{\lambda_f}$ identifies two arguments θ and θ' if and only if the external rays $R_f(\theta)$ and $R_f(\theta')$ land at a common point. In this case, L is an infinite λ-unlinked if and only if there exists a bounded Fatou component V such that L is the set formed by the arguments of all the irrational rays landing at ∂V. Finite unlinked classes are irrational equivalence classes of $\widehat{\lambda_f}$ and correspond to external rays landing at a common point, with infinite forward orbit.

The main properties of λ-unlinked classes and their fiber is summarized in the proposition below. Before we state the proposition let us agree that if $X, Y \subset \mathbb{C}$ and $f : X \to Y$ is a surjective map defined and holomorphic on a neighborhood of X, then we say that the *degree of $f : X \to Y$* is $\delta \geq 1$ if every point in Y has δ preimages on X, counting multiplicities.

Proposition 3.7. *Consider a d-invariant rational lamination λ and $f \in \mathcal{C}(\lambda)$. Let L be a λ-unlinked class. Then*

(i) *$m_d(L)$ is a λ-unlinked class.*
(ii) *$f(K_f(L)) = K_f(m_d(L))$.*

(iii) *If L is finite, then*
  (a) *L is a $\hat{\lambda}$-class. In particular, L is wandering ($m_d^n(L) \neq m_d^m(L)$ for any $n \neq m$).*
  (b) *$m_d : L \to m_d(L)$ is $\delta = \delta(L)$-to-one for some $\delta \geq 1$.*
  (c) *The degree of $f : K_f(L) \to K_f(m_d(L))$ is well defined and exactly $\delta(L)$.*
(iv) *If L is infinite, then*
  (a) *L is eventually periodic (i.e., there exists $n > 0$ and $\ell \geq 0$ such that $m_d^{n+\ell}(L) = m_d^{\ell}(L)$).*
  (b) *If $m_d^n(L) = L$, then $\delta^n(L) > 1$, where $\delta^n(L) = \prod_{k=0}^{n-1} \delta(m_d^k(L))$.*
  (c) *$\overline{L}/\hat{\lambda}$ is homeomorphic to $\mathbb{R}/\mathbb{Z}$. Moreover, if (θ, θ') is a connected component of $\mathbb{R}/\mathbb{Z} \setminus \overline{L}$, then θ and θ' are in $\mathbb{Q}/\mathbb{Z}$ and λ-equivalent. Furthermore, $(m_d(\theta), m_d(\theta'))$ is a connected component of $\overline{m_d(L)}$ or $m_d(\theta) = m_d(\theta')$.*
  (d) *$m_d : \overline{L} \to \overline{m_d(L)}$ induces a map $\tilde{m}_d : \overline{L}/\hat{\lambda} \to \overline{m_d(L)}/\hat{\lambda}$, which is conjugate to the δ-fold covering map $m_\delta : \mathbb{R}/\mathbb{Z} \to \mathbb{R}/\mathbb{Z}$ for some $\delta = \delta(L) \geq 1$.*
  (e) *Assume further that for any λ_f-class A and any λ-unlinked class L' such that $A \cap \overline{L} = \emptyset$ and $m_d(L') = m_d(L)$ we have $A \cap \overline{L'} = \emptyset$. Then the degree of $f : K_f(L) \to K_f(m_d(L))$ is exactly $\delta(L)$.*

The reader may find the proof of the above proposition at the end of this section.

The following examples show that the further assumptions in property (iv)(e) are necessary.

Example 3.8. 1. Consider a family $P_a(z) = az^3 - (a+1)z^2 + 1$ of cubic polynomials having a period two superattractive cycle $\{0, 1\}$. Let $f_0 = P_{a_0}$ and $f_1 = P_{a_1}$ where $a_0 = -1/4$ and $a_1 = \frac{11-3\sqrt{17}}{4}$ (see Figure 4). The critical point $c_a = \frac{2(a+1)}{3a}$ is mapped to 0 for f_0 and for f_1, it is mapped to the fixed point which is the intersection of the closures of the immediate basins of 0 and 1.

Although $\lambda_0 = \lambda_{f_0}$ is admissible for f_1, the polynomial f_1 is not λ_0-renormalizable. In fact, let L (resp. L') be the λ_0-unlinked class whose λ_0-fiber for f_0 contains 0 (resp. c_{a_0}). Then, we have $K_{f_1}(L) \cap K_{f_1}(L') = \{c_{a_1}\}$. Therefore, $K_{f_1}(L)$ contains both of the critical points and any proper extension of $f_1 : K_{f_1}(L) \to K_{f_1}(L')$ to a disk would have degree 3.

The assumptions of (iv)(e) do not hold. In fact, let L'' be the λ-unlinked class distinct from L' such that $m_d(L) = m_d(L'')$. Note that $\overline{L''} \cap \overline{L'} \neq \emptyset$. Also, observe that $A_0 = \overline{L} \cap \overline{L'}$ is a λ_0-class. The λ_{f_1}-class

$$A = A_0 \cup (\overline{L''} \cap \overline{L'}),$$

formed by the the angles landing at c_{a_1}, intersects both $\overline{L}$ and $\overline{L''}$.

2. Let us now illustrate a more complicated situation in which the assumptions of (iv)(e) fail to hold. Namely, the assumption fails for a λ_0-class A intersecting the closure of a λ_0-unlinked class L. However, L is the unique critical λ_0-unlinked class whose closure intersects A.

Let $g_0(z) = 2.959...z^3 + 2.418...z^5 + 0.4590...z^7$ and $g_1(z) = -1.258...z + 0.06411...z^3 + 0.4047...z^5 + 0.08182...z^7$ (see Figure 5). They are odd polynomials of degree 7. The critical points of g_0 are 0 (double critical point), $\pm c_1 = \pm i$ and $\pm c_2 = \pm 1.662...i$, which satisfy $g_0(0) = 0$, $g_0(\pm c_1) = g_0(\mp c_2) = \mp c_1$. In particular, g_0 is post-critically finite and hyperbolic. For g_1, the critical points are $\pm c_1' = \pm i$, $\pm c_2' = \pm 1.793...i$ and $\pm c_3 = \pm 0.8265...$ and we have $g_1(\pm c_1) = \mp c_1$, $g_1(\pm c_2) = 0$ and $g_2(\pm c_3) = \mp c_3$. Furthermore, we have $\lambda_{g_1} \supset \lambda_{g_0}$. Let L be the λ_{g_0}-unlinked classes such that $c_1 \in K_{g_0}(L)$ for $i = 1, 2$. Then $K_{g_1}(L)$ contains both c_1 and c_2, so similar to the previous example, this implies the conclusion of (iv)(e) does not hold.



FIGURE 4. Yoccoz puzzles and the rational laminations for the monic centered polynomials affinely conjugate to $f_a(z) = az^3 - (a+1)z^2 + 1$ for $a = -1/4$ and $a = \frac{11-3\sqrt{17}}{4} = -0.3423...$ Dots indicate critical points.

Observe that there are two bifurcations from g_0 to g_1: The first one is that period two points collapse at the origin and new landing relations indicated by the light gray regions in the picture of the rational lamination for g_1 appear. The second is that the critical points $\pm c_2'$ hit the origin, the corresponding landing relations are indicated by the dark gray regions.

Below we combinatorially and abstractly describe the "location" and "orbit" of the critical points.

Definition 3.9 (Critical elements and orbits)**.** Let λ be a d-invariant rational lamination and $\hat{\lambda}$ its real extension (Definition 3.6). If A is a $\hat{\lambda}$-class, then $\delta(A)$ denotes the degree of $m_d : A \to m_d(A)$. If A is a $\hat{\lambda}$-class with $\delta(A) > 1$, then we say that A is a *Julia critical element* of λ.



Figure 5. Yoccoz puzzles and the rational laminations for the monic centered polynomials affinely conjugate to $g_0(z) = 2.959...z^3 + 2.418...z^5+0.4590...z^7$ and $g_1(z) = -1.258...z+0.06411...z^3+0.4047...z^5+0.08182...z^7$. Dots indicate critical points.

If L is an infinite unlinked λ-class, we say that $\delta(L)$ is the *degree of L* where $\delta(L)$ is the number obtained in Proposition 3.7 (iv)(d). If $\delta(L) > 1$, then we say that L is a *Fatou critical element* of λ.

We denote by $\mathrm{Crit}(\lambda)$ the collection formed by all the critical elements for λ. The *post-critical set* and *critical orbits of λ* are:

$$PC(\lambda) = \{m_d^n(C);\ C \in \mathrm{Crit}(\lambda), n > 0\}$$

$$CO(\lambda) = \mathrm{Crit}(\lambda) \cup PC(\lambda)$$

It follows that

$$d = 1 + \sum_{C\in\mathrm{Crit}(\lambda)} (\delta(C) - 1)$$

[Ki, Lemma 4.10].

Definition 3.10 (Hyperbolic and post-critically finite laminations)**.** An invariant rational lamination λ is *hyperbolic* if it has no Julia critical element. An invariant rational lamination λ is *post-critically finite* if every Julia critical element is contained in $\mathbb{Q}/\mathbb{Z}$.

It follows that a hyperbolic invariant rational lamination is the rational lamination of post-critically finite hyperbolic polynomial. Also, a post-critically finite invariant rational lamination is the rational lamination of a post-critically finite polynomial [Ki].

3.3. **Renormalizations.**

Definition 3.11 (Mapping schema associated to λ_0)**.** Consider an invariant rational lamination λ_0.

- Let $|T(\lambda_0)|$ be the collection formed by the Fatou critical elements v of λ_0.
- Let $\sigma_{\lambda_0} : |T(\lambda_0)| \to |T(\lambda_0)|$ be the map such that $\sigma_{\lambda_0}(v) = v'$ if $m_d^{\ell_v}(v) = v'$, for some $\ell_v \geq 0$, and $m_d^k(v) \notin |T(\lambda_0)|$, for all $0 < k < \ell_v$. We say that ℓ_v is the *return time* of v.
- Finally, let $\delta_{\lambda_0} : |T(\lambda_0)| \to \mathbb{N}$ be the corresponding degree map. That is, $\delta_{\lambda_0}(v) = \delta(v)$.

We call

$$T(\lambda_0) = (|T(\lambda_0)|, \sigma_{\lambda_0}, \delta_{\lambda_0}).$$

the *reduced mapping schema of* λ_0.

Definition 3.12 (Renormalizable and renormalization)**.** We say $f \in C(\lambda_0)$ is λ_0*-renormalizable* if, for every $v \in |T(\lambda_0)|$ there exist topological disks U'_v and U_v such that

$$g = (f^{\ell_v} : U'_v \to U_{\sigma_{\lambda_0}(v)})_{v\in|T(\lambda_0)|}$$

is a polynomial-like map over $T(\lambda_0)$ with fiberwise connected filled Julia set

$$K(g) = \bigcup_{v\in|T(\lambda_0)|} \{v\} \times K_f(v).$$

We call g a λ_0*-renormalization* of f. We denote by $\mathcal{R}(\lambda_0)$ the subset of $C(\lambda_0)$ formed by all the λ_0-renormalizable polynomials.

The above definition is stronger than that in the outline (Section 1), but they are in fact equivalent.

Proposition 3.13. *Let $f \in C(\lambda_0)$. If there exist topological disks U'_v and U_v for each $v \in |T(\lambda_0)|$ such that $K_v(f) \subset U'_v$ and $g = (f^{\ell_v} : U'_v \to U_{\sigma_{\lambda_0}(v)})_{v\in|T(\lambda_0)|}$ is a polynomial-like map over $T(\lambda_0)$, then g is a λ_0-renormalization of f.*

Proof. It suffices to show $K_f(v) = K(g, v)$. By construction, we have $K_f(v) \subset K(g, v)$. Since v is an infinite λ_0-unlinked class, $K_f(v)$ is a full continuum. For v periodic of period n by m_d, let $U''_v = f^{-n}(U_v) \cap U'_v$. Then $f^n : U''_v \to U_v$ is a polynomial-like map and its filled Julia set is $K(g, v)$). Hence $K(g, v)$ is the smallest full continuum in U''_v completely invariant by $f^n : U''_v \to U_v$. Therefore, $K_f(v) \supset K(g, v)$. For preperiodic $v \in |T(\lambda_0)|$, we need only take a backward image of $K(g, v')$ where $v' = \sigma^k(v)$ is periodic. □

In order to extract polynomial-like maps with an external marking it is convenient to "internally mark" the invariant rational lamination λ_0.

Definition 3.14 (Internally angled)**.** An *internally angled invariant rational lamination* λ_0 is a pair $(\lambda_0, (\alpha_v : \overline{v} \to \mathbb{R}/\mathbb{Z})_{v\in|T(\lambda_0)|})$ such that

(i) α_v induces a homeomorphism from $\overline{v}/\lambda_0$ to $\mathbb{R}/\mathbb{Z}$.

(ii) $\alpha_{\sigma_{\lambda_0}(v)}(d\theta) = \delta_{\lambda_0}(v)\alpha_v(\theta)$ for $\theta \in \overline{v}$.

We call $(\alpha_v)_{v\in|T(\lambda_0)|}$ an *internal angle system*.

The existence of internal angle system is guaranteed by Proposition 3.7. An internally angled invariant rational lamination determines an external marking the λ_0-renormalization of every $f \in \mathcal{R}(\lambda_0)$.

Definition 3.15 (Induced external marking)**.** Let $(\lambda_0, (\alpha_v : \overline{v} \to \mathbb{R}/\mathbb{Z})_{v\in|T(\lambda_0)|})$ be an internally angled invariant rational lamination. For each $v \in |T(\lambda_0)|$ choose an argument θ_v in $\alpha_v^{-1}(0)$. Given a λ_0-renormalization g of a polynomial $f \in \mathcal{R}(\lambda_0)$, let Γ_v be the access with representative $R_f(\theta_v) \cap U'_v$. We say that $\Gamma = (\Gamma_v)_{v\in|T(\lambda_0)|}$ is the *external marking of g determined by the internal angle system* $(\alpha_v : \overline{v} \to \mathbb{R}/\mathbb{Z})_{v\in|T(\lambda_0)|}$.

By Lindelöf Theorem, the external marking of g determined by an internal angle system is independent of the choices involved in the definition above.

Lemma 3.16. *Let λ_0 be an invariant rational lamination. Let g be a λ_0-renormalization of $f \in \mathcal{R}(\lambda_0)$ and Γ an external marking of g. Then there exists an internal angle system α such that the external marking determined by α is exactly* Γ.

Proof. Let $\beta = (\beta_v)$ be an internal angle system for λ_0. Consider the straightening of g that maps the external marking induced by β onto the standard marking of $P = \chi_{\lambda_0}(g)$. It follows that Γ is mapped onto an external marking of P determined by a collection of external rays $(R_P(\theta_v, v))$. Now the internal angle system $\alpha = (\alpha_v = \beta_v + \theta_v)$ is such that the induced external marking Γ_α of g maps onto the external marking of P determined by $(R_P(\theta_v, v))$. Therefore, $\Gamma = \Gamma_\alpha$. □

3.4. **Proof of Proposition 3.7.** Most of this proposition is already proved in [Ki, Section 4] and [In3, Section 3]. In fact, it only remains to establish properties (ii), (iii)(c) and (iv)(e).

It is convenient to consider a (countable) set M contained in $\lambda_0 \subset \mathbb{Q}/\mathbb{Z} \times \mathbb{Q}/\mathbb{Z}$ formed by the "image" of non-trivial relations. More precisely,

$$M = \{(d\theta, d\theta') \in \lambda_0; \theta \neq \theta', (\theta, \theta') \in \lambda_0\}.$$

Given a λ_0-unlinked class L, the invariance of λ_0 implies that:

$$K_f(m_d(L)) = K(f) \cap \bigcap_{(\theta,\theta')\in M} \overline{\mathrm{Sector}(\theta, \theta'; m_d(L))},$$

$$K_f(L) = K(f) \cap \bigcap_{(\theta,\theta')\in M} \overline{S'((\theta, \theta'))},$$

where

$$S'((\theta, \theta')) = \bigcap_{(\theta_1,\theta_2)\in\lambda_0,\ d\theta_1=\theta,\ d\theta_2=\theta'} \mathrm{Sector}(\theta_1, \theta_2; L)$$

is the intersection of all the components of $f^{-1}(\mathrm{Sector}(\theta, \theta'; m_d(L)))$ containing $R_f(\theta)$ for some $\theta \in L$.

Now (ii) follows, after countable intersection of the equation,

$$f\left(\overline{S'(\theta, \theta')}\right) = \overline{\mathrm{Sector}(\theta, \theta'; m_d(L))},$$

which holds for all $(\theta, \theta') \in M$.

To prove (iii)(c), we assume that L is finite and recall that, by (iii)(a), $L \subset (\mathbb{R} \setminus \mathbb{Q})/\mathbb{Z}$. Hence $K_f(L) \subset S'((\theta, \theta'))$ for all $(\theta, \theta') \in M$. Since, $f : S'((\theta, \theta')) \to \mathrm{Sector}(\theta, \theta'; m_d(L))$ is a proper map for all $(\theta, \theta') \in M$, property (iii)(c) follows.

Now assume L is infinite and satisfies the assumption of (iv)(e). Let $M_L = \{(\theta, \theta') \in M;\ \theta, \theta' \in \overline{m_d(L)}\}$ and for each $(\theta, \theta') \in M_L$, let

$$M'_L(\theta, \theta') = \{(\theta_1, \theta_2) \in M;\ d\theta_1 = \theta,\ d\theta_2 = \theta',\ \theta_j \in \overline{L},\ (j = 1, 2)\}.$$

Then we have

$$K_f(m_d(L)) = K(f) \cap \bigcap_{(\theta,\theta') \in M_L} \overline{\mathrm{Sector}(\theta, \theta'; m_d(L))},$$
$$K_f(L) = K(f) \cap \bigcap_{(\theta,\theta') \in M_L} \overline{S''(\theta, \theta')},$$

where

$$S''(\theta, \theta') = \bigcap_{(\theta,\theta') \in M'_L} \mathrm{Sector}(\theta, \theta'; L).$$

The assumption guarantees that, for all $(\theta, \theta') \in M_L$, each component K of $f^{-1}(K_f(m_d(L)))$ is either contained in or disjoint from $\overline{S''(\theta, \theta')}$. Hence, for all $(\theta, \theta') \in M_L$,

$$f : f^{-1}(K_f(m_d(L))) \cap \overline{S''(\theta, \theta')} \to K_f(m_d(L))$$

has degree at least $\delta(L)$. After taking a countable intersection, property (iv)(e) follows. □

4. YOCCOZ PUZZLES

First, we introduce combinatorial Yoccoz puzzles in terms of laminations, then we define Yoccoz puzzles corresponding to a combinatorial Yoccoz puzzle. See [In3] for more details.

Throughout this section we fix an integer $d \geq 2$.

4.1. **Combinatorial Yoccoz Puzzle.**

Definition 4.1. A *combinatorial Yoccoz puzzle* $\Lambda = (\Lambda_k)_{k \geq 0}$ is a sequence of rational laminations with finite support such that:

(i) If $\theta \in \mathrm{supp}(\Lambda_0)$, then θ is periodic under m_d.
(ii) If A is a Λ_0-class, then $m_d(A)$ is a Λ_0-class and $m_d : A \to m_d(A)$ is consecutive preserving.
(iii) $(m_d)_* \Lambda_{k+1} = \Lambda_k$ for any $k \geq 0$.
(iv) If A is a non-trivial λ_{k+1}-class, then $m_d(A)$ is also a non-trivial Λ_k-class.

We say that $\mathrm{supp}(\Lambda) = \bigcup_k \mathrm{supp}(\Lambda_k)$ is the *support of* Λ.

A Λ_k-unlinked class is a *combinatorial puzzle piece of depth k*. The collection of all depth k combinatorial puzzle pieces is denoted by $\mathcal{L}_k(\Lambda)$.

We say that a combinatorial Yoccoz puzzle Λ is *admissible for a lamination λ* if $\Lambda_k \subset \lambda$ for all $k \geq 0$. We say Λ is *admissible for a polynomial $f \in \mathcal{C}(d)$* if Λ is admissible for λ_f.

We state, without proof, the following rather straightforward properties of combinatorial puzzles.

Proposition 4.2. *Let $\Lambda = (\Lambda_k)_k$ be a combinatorial Yoccoz puzzle. Then*

(i) $\{\mathrm{supp}(\Lambda_k)\}_k$ *forms an increasing sequence in $\mathbb{Q}/\mathbb{Z}$ and* $\mathrm{supp}(\Lambda)$ *is completely invariant under m_d.*
(ii) *For all $k \geq 0$, if L is combinatorial piece of a depth $k + 1$, then there exists a unique combinatorial piece L' of depth k such that $L \subset L'$.*

(iii) *For all* $k \geq 0$*, if* L' *is a depth* $k+1$ *combinatorial piece of a depth* $k+1$*, then* $m_d(L')$ *is a combinatorial piece of depth* k *and* $m_d : L' \to m_d(L)$ *is* δ*-to-one for some* $\delta \geq 1$*.*

(iv) *Let* L *be a combinatorial piece of depth* k*, for some* $k \geq 0$*. If* $\theta \in \partial L$*, then for all* $k' > k$*, there exists a unique combinatorial piece* L' *of depth* k' *such that* $L' \subset L$ *and* $\theta \in \partial L'$*.*

(v) *Let* $L_1, \dots, L_n$ *be a complete list (without repetitions) of the combinatorial puzzle pieces of depth* k*. Let* $C_1, \dots, C_m$ *be a complete list (without repetitions) of the* Λ_k*-critical classes. Then,*

$$\sum_{i=1}^{n}(\delta(L_i) - 1) + \sum_{j=1}^{m}(\delta(C_j) - 1) = d - 1,$$

where, for $A = L_i$ *or* C_j*, the number* $\delta(A)$ *denotes the degree of* $m_d : A \to m_d(A)$*.*

For a combinatorial puzzle piece L, we denote by $\delta(L)$ the degree of $m_d : L \to m_d(L)$.

4.2. **Yoccoz Puzzle.** Let f be a monic centered polynomial with connected Julia set and associated Böttcher map $\varphi : \mathbb{C} \setminus K(f) \to \mathbb{C} \setminus \overline{\Delta}$ where $\Delta = \{|z| \leq 1\}$. For $r > 0$, the Jordan curve

$$E_f(r) = \varphi^{-1}(\{z \in \mathbb{C} \setminus \overline{\Delta};\, |z| = \exp(r)\})$$

is called the *equipotential of level* r. The open topological disk bounded by $E_f(r)$ will be denoted by $D_f(r)$.

Fix $r > 0$. Let $\Lambda = (\Lambda_k)$ be an admissible combinatorial Yoccoz puzzle for $f \in \mathcal{C}(d)$. We define the *depth* k *Yoccoz puzzle for* (f, Λ) as follows. Given a combinatorial puzzle piece L of depth k, let

$$P_k(f, L) = \overline{D_f(r/d^k)} \cap \bigcap_{\theta \sim_{\Lambda_k} \theta', \theta \neq \theta'} \overline{\mathrm{Sector}(\theta, \theta'; L)}$$

be the *puzzle piece of depth* k *for* (f, Λ) *associated to* L. We omit k and/or f when clear by context. Recall that we denote the collection of combinatorial puzzle pieces of depth $k \geq 0$ by $\mathcal{L}_k(\Lambda)$. Similarly we let

$$\mathcal{P}_k(f, \Lambda) = \{P(L);\ L \in \mathcal{L}_k(\Lambda)\}$$

be the collection of all puzzle pieces of depth k. That is, the *Yoccoz puzzle of depth* k *for* (f, Λ).

Remark 4.3. The interior of a puzzle piece is not connected in general. A "degenerate" puzzle appears when two distinct Λ_k-classes A, A' are contained in the same λ-class. Therefore, our definition of puzzle piece does not coincide with the usual one (i.e., the closure of a bounded component of the complement of a suitable graph constructed with equipotentials and external rays).

Example 4.4. 1. Let $c = -0.1010... + 0.9562...i$ and $f(z) = z^2 + c$, so that $\alpha = f^4(c)$ is the alpha fixed point (i.e., the fixed point which is not a landing point of $R_f(0)$). Figure 6 shows its Julia set with its Yoccoz puzzles and (a part of) its lamination. The rational lamination λ_f of f properly contains the rational lamination of Douady's rabbit (say λ_0). In fact, the critical value c has the same landing angles of the alpha fixed point, i.e., $1/7, 2/7, 4/7$. Thus the critical point 0 has six landing angles, consisting of two λ_0-equivalence classes.

The lightgray region in the right picture indicates the landing relation for the critical point 0, which is not contained in the lamination of Douady's rabbit. The Yoccoz puzzles



FIGURE 6. The Julia set and a "degenerate" puzzle for $z^2 + c$ with $c = -0.1010... + 0.9562...i$.

for f, determined by the lamination of Douady's rabbit (namely, let E_k be the set of landing angles for $f^{-k}(\alpha)$ and $\Lambda_k = \lambda|_{E_k}$), is such that the puzzle piece of depth 4 containing 0 has two interior components.

2. Let $f_0(z) = z^3 + az^2$ where $a = 1.502... - 0.7790...i$ and let $f_1(z) = z^3 - 3/4z - \sqrt{7}i/4$ (Figure 7). There are two superattractive cycles of period one and two for f_0, and f_1 has two superattractive cycles of period two. Observe that $\lambda_{f_0} \subset \lambda_{f_1}$. Every non-trivial class of λ_{f_0} eventually maps onto the fixed class formed by the angles $1/4, 5/8$. As above, we consider the Yoccoz puzzles (Λ_k) determined by λ_{f_0}. The puzzle piece for f_1 of any depth corresponding to the fixed superattractive basin for f_0 has disconnected interior (and the number of components increases as depth increases).

3. Similarly, the polynomials f_1 and g_1 in Example 3.8 are also examples having degenerate puzzle pieces.

Proposition 4.5. *Let $\Lambda = (\Lambda_k)$ be an admissible combinatorial Yoccoz puzzle for $f \in \mathcal{C}(d)$. For all $k \geq 0$, let*

$$\Gamma_k = E_f(r/d^k) \cup \bigcup_{\theta \in \operatorname{supp}(\Lambda_k)} \overline{R_f(\theta)}.$$

Then the following statements hold:

(i) *If L is a combinatorial puzzle piece, then $P(L)$ is compact, connected and full. Moreover, the interior of $P(L)$ is the union of all the bounded components of $\mathbb{C} \setminus \Gamma_k$ which intersect $R_f(\theta)$ for some $\theta \in L$.*

(ii) *$\mathcal{P}_k(f, \Lambda)$ is a partition of $\overline{D_f(r/d^k)}$, i.e., depth k puzzle pieces have mutually disjoint interior and*

$$\overline{D_f(r/d^k)} = \bigcup_{P \in \mathcal{P}_k(f,\Lambda)} P.$$

(iii) *If combinatorial puzzle pieces L and L' satisfy $L \subset L'$, then $P(L) \subset P(L')$.*

(iv) *If L is a combinatorial puzzle piece of depth $k \geq 1$, then $f(P(L)) = P(m_d(L))$ and $f : \operatorname{int} P(L) \to \operatorname{int} P(m_d(L))$ is a proper map of degree $\delta(L)$.*



FIGURE 7. Yoccoz puzzles and the rational laminations for $f_0(z) = z^3+az^2$ ($a = 1.502... - 0.7790...i$) and $f_1(z) = z^3 - 3/4z - \sqrt{7}i/4$.

For the proof, see [In3, Proposition 4.1]. This proposition implies that our puzzles have similar properties as the usual Yoccoz puzzles (defined by the closures of bounded components of $\mathbb{C} \setminus \Gamma_k$) and moreover, their properties can be described only in terms of Λ.

4.3. **From combinatorial puzzles to laminations.** Combinatorial Yoccoz puzzles are particularly suited to study the smallest invariant rational lamination for which they are admissible.

Definition 4.6. Let $\Lambda = (\Lambda_k)$ be a combinatorial Yoccoz puzzle. The *rational lamination* $\lambda(\Lambda)$ *generated by* Λ is the smallest equivalence relation in $\mathbb{Q}/\mathbb{Z}$ containing the closure of $\bigcup_k \Lambda_k$ in $\mathbb{Q}/\mathbb{Z} \times \mathbb{Q}/\mathbb{Z}$.

The previous definition is justified with the following result.

Proposition 4.7. *For a combinatorial Yoccoz puzzle* $\Lambda = (\Lambda_k)_{k\geq 0}$, *the rational lamination* $\lambda(\Lambda)$ *generated by* Λ *is an invariant rational lamination.*

Proof. First, it is clear that $\lambda(\Lambda)$-classes are pairwise unlinked. Applying the argument for the claim in the proof of [Ki, Lemma 4.7] to our case, we have the following:

Claim. If $\theta, \theta' \in \mathbb{Q}/\mathbb{Z}$ are $\overline{\lambda}$-related but not λ-equivalent, and if θ is periodic by d, then θ' is periodic of the same period.

If A is a $\lambda(\Lambda)$-class, then $m_d^n(A)$ contains a periodic angle under iterations of m_d. Therefore, $m_d^n(A)$ (hence A) is a finite set since, by the claim above, $m_d^n(A)$ consists of periodic angles of the same period.

Now let us prove that $\lambda(\Lambda)$ is closed. Assume that, for all $n \geq 0$, θ_n, θ'_n are $\lambda(\Lambda)$-equivalent arguments such that $\theta_n \to \theta$ and $\theta'_n \to \theta'$, as $n \to \infty$. Then, since $\lambda(\Lambda)$-classes are pairwise unlinked, we may assume that these sequences are monotone. That is, $\theta_n \searrow \theta$ and $\theta'_n \nearrow \theta'$. Moreover, we may also assume that, for all n, the $\lambda(\Lambda)$-class containing θ_n is contained in $[\theta_n, \theta_{n-1}) \cup (\theta'_{n-1}, \theta'_n]$. It follows that there exists some $\tilde{\theta}_n \in [\theta_n, \theta_{n-1})$ and $\tilde{\theta}'_n \in (\theta_{n-1}, \theta_n]$ such that $\tilde{\theta}_n, \tilde{\theta}'_n$ are λ-equivalent. Hence, θ, θ' are $\overline{\lambda}$-related and, therefore, $\lambda(\Lambda)$-equivalent.

It remains to prove that $(m_d)_* \lambda(\Lambda) = \lambda(\Lambda)$. It is clear that for a $\lambda(\Lambda)$-class A, $m_d(A)$ is contained in a $\lambda(\Lambda)$-class B. Assume there exists $\theta'_0 \in B \setminus m_d(A)$. Take $\theta_1, \theta_2 \in A$ such that $(d\theta_1, d\theta_2)$ is the component of $\mathbb{R}/\mathbb{Z} \setminus m_d(A)$ containing θ'_0. We may assume θ'_0 and $d\theta_1$ are $\overline{\lambda}$-related (because either $\theta_1 \sim_{\lambda(\Lambda)} \min[(d\theta_1, d\theta_2) \cap B]$ or $\theta_2 \sim_{\lambda(\Lambda)} \max[(d\theta_1, d\theta_2) \cap B]$ holds). Then there exists $\theta'_{0,n} \nearrow \theta'_0$ and $\theta'_{1,n} \searrow d\theta_1$ such that $\theta'_{0,n}$ and $\theta'_{1,n}$ are λ-equivalent. Let $\theta_{1,n} \searrow \theta_1$ be such that $d\theta_{1,n} = \theta'_{1,n}$. Then there exists $\theta_{0,n}$ such that $d\theta_{0,n} = \theta'_{0,n}$ and $\theta_{0,n}, \theta_{1,n}$ are λ-equivalent. Since $\lambda(\Lambda)$-classes are pairwise unlinked, $\theta_{0,n}, \theta_{1,n} \subset (\theta_1, \theta_2)$. Passing to a subsequence, we may assume $\theta_{0,n} \to \theta_0$ and $d\theta_0 = \theta'_0$. Then since θ_0 and θ_1 are $\overline{\lambda}$-related and $\theta_0 \in [\theta_1, \theta_2]$, either $\theta_0 = \theta_1$ or $\theta_0 = \theta_2$, it is a contradiction. Hence $B = m_d(A)$ is a $\lambda(\Lambda)$-class. This argument also shows that $m_d : A \to m_d(A)$ is consecutive preserving. □

Definition 4.8. Let $\Lambda = (\Lambda_k)$ be an admissible combinatorial Yoccoz puzzle for a d-invariant rational lamination λ. We say Λ is a *generator of* λ if there exists $k \geq 0$ such that, for any combinatorial piece L of depth k which contains a λ-unlinked class in the forward orbit of a Fatou critical element of λ, we have that $\delta(L) = \delta(v)$. We call such k a *separation depth* for λ.

The next proposition shows that a generator actually "generates" a substantial part of the corresponding rational lamination.

Proposition 4.9. *Let $\Lambda = (\Lambda_k)_{k \geq 0}$ be a generator of an invariant rational lamination λ. If L is a Fatou critical element of λ, then L is a Fatou critical element of $\lambda(\Lambda)$.*

Proof. Let $\lambda' = \lambda(\Lambda)$ and observe that $\lambda' \subset \lambda$. Therefore, given any infinite λ-unlinked class L, there exists a unique λ'-unlinked class L' such that $L' \supset L$.

Let L be a Fatou critical element of λ and L' the λ-unlinked class containing L. We claim that $L' = L$. For this purpose, let k be a separation depth for λ. Denote by L_k the combinatorial puzzle piece of depth k containing L'. Thus, $\delta(L_k) = \delta(L') = \delta(L)$. Now assume that L is periodic under m_d, say of period p. It follows that $m_d^p : L' \to L'$ and $m_d^p : L \to L$ have the same degree. Moreover, under both of these maps, the grand orbits of a given $\theta \in L$ coincide. Since such a grand orbit is dense in L and in L', we conclude that $L = L'$. In the case that L is preperiodic, then there exists $\ell \geq 1$ such that $m_d^\ell(L) = m_d^\ell(L')$ is periodic and the degree of $m_d^\ell : L \to m_d^\ell(L)$ also coincides with the degree of $m_d^\ell : L' \to m_d^\ell(L')$. Hence, $L = L'$. □

Lemma 4.10. *If λ be an invariant rational lamination, then there exists a generator of λ.*

Proof. Given v a λ-unlinked class in the forward orbit of a Fatou critical element of λ, according to Proposition 3.7 (iv), for every critical element $C \neq v$ of λ there exists a λ-class A_C such that the connected component of $\mathbb{R}/\mathbb{Z} \setminus A_C$ that contains v is disjoint from C. Hence, there exists a collection $A_1, \dots, A_n$ of λ-classes such that the following holds. If v is a λ-unlinked class in the forward orbit of a Fatou critical element of λ and C is a critical element of λ, then the connected component of $\mathbb{R}/\mathbb{Z} \setminus A_j$ that contains v is disjoint from C, for some $j = 1, \dots, n$. Let E_0 be the periodic arguments in the m_d-forward orbit of $A_1 \cup \cdots \cup A_n$. For $k \geq 0$, let $E_k = m^{-k}(E_0)$ and Λ_k be the restriction of λ to $E_k \times E_k$. It follows that the puzzle $\Lambda = (\Lambda_k)_{k \geq 0}$ is a generator for λ. □

5. Injectivity of straightening maps

This section is devoted to establish that straightening maps are injective. That is, we prove Theorem B. The proof relies on the following main result.

Key Lemma 5.1. *Let λ_0 be a post-critically finite d-invariant rational lamination with reduced schema $T(\lambda_0)$. If $f \in \mathcal{C}(\lambda_0)$, then the set*

$$F = K(f) \setminus \bigcup_{n \geq 0} \bigcup_{v \in |T(\lambda_0)|} f^{-n}(K_f(v))$$

has zero area.

Section 5.2 contains the proof of this lemma. In Section 5.1 we prove Theorem B assuming the Key Lemma.

Remark 5.2. Epstein and Yampolsky [EY] first proved this key lemma for cubic polynomials constructed by intertwining surgery of two quadratic polynomials. Haïssinsky [Ha3] also claimed the key lemma for a general intertwining surgery, but his proof contains a gap. (He claimed that the set denoted by F_1, in the proof below, is empty. However, F_1 is always nonempty in the case of intertwining surgery)

Remark 5.3. Theorem B often implies that the polynomial obtained through a quasiconformal construction from a collection of polynomials is uniquely determined by the original polynomials, up to affine conjugacy. For example, consider the case of intertwining surgery by Epstein and Yampolsky [EY]. Starting with two quadratic polynomials with connected Julia sets they construct a cubic polynomial (called an intertwining) having two quadratic-like restrictions whose Julia sets intersect at a repelling fixed point. The resulting cubic polynomial only depends on the choice of initial quadratic polynomials and not on the choices made throughout the surgery.

The first author [In2] used the intertwining construction to obtain a polynomial having a capture type renormalization. For such a construction, Theorem B can be applied to show that the resulting polynomial is uniquely determined by the initial data, up to affine conjugacy. Haïssinsky [Ha3] also constructed intertwinings at parabolic fixed points which produce a non-renormalizable polynomial. However, the Key Lemma 5.1 still holds and, therefore, the resulting polynomial of the construction is uniquely determined by the corresponding initial data, up to affine conjugacy.

5.1. Proof of Theorem B.

First, we assume the key lemma and prove Theorem B.

Let λ_0 be a post-critically finite d-invariant rational lamination and let α be an internal angle system for λ_0. Consider two renormalizable polynomials $f_1, f_2 \in \mathcal{R}(\lambda_0)$ such that

$\chi_{\lambda_0}(f_1) = \chi_{\lambda_0}(f_2)$, where χ_{λ_0} is the corresponding straightening map. We must show that $f_1 = f_2$.

For the purpose of this proof it is better to extract "polynomial-like maps" g_1, g_2 over the *unreduced mapping schema* for λ_0. More precisely, let $|T^U(\lambda_0)|$ be the (forward) orbit of the Fatou critical elements of λ_0 and, for all $v \in |T^U(\lambda_0)|$, let $\delta(v)$ be the degree of $m_d : v \to m_d(v)$. It follows that $T^U(\lambda_0) = (|T^U(\lambda_0)|, m_d, \delta)$ is a mapping schema. If $f \in \mathcal{R}(\lambda_0)$, then there exists domains $U'_v \Subset U_v$ such that $g_v = f : U'_v \to U_{m_d(v)}$ is a proper map of degree $\delta(v)$, for all $v \in |T^U(\lambda_0)|$. Moreover, these domains may be chosen so that the forward orbit of $(v, z) \in |T^U(\lambda_0)| \times \mathbb{C}$ under $(v, z) \mapsto (m_d(v), f(z))$ is well defined if and only if $z \in K_v(f)$. We say that $g = (f : U'_v \to U_{m_d(v)})_{v \in |T^U(\lambda_0)|}$ is an *unreduced λ_0-renormalization* of f.

Now let $g_j = (f_j : U'_{j,v} \to U_{j,m_d(v)})_{v \in |T^U(\lambda_0)|}$ be an unreduced λ_0-renormalization of f_j, for $j = 1, 2$. Given a hybrid conjugacy between the λ_0-renormalizations of f_1 and f_2, that respects the external markings, determined by the internal angle system α, it is not difficult to extend it to a hybrid conjugacy ψ between g_1 and g_2. That is, $\psi = (\psi_v : U_{1,v} \to U_{2,v})_{v \in |T^U(\lambda_0)|}$ is such that, for all $v \in |T^U(\lambda_0)|$:

$$\frac{\partial \psi_v}{\partial \overline{z}}(z) \equiv 0,$$

for almost every $z \in K_v(f_1)$ and,

$$f_2 \circ \psi_v(z) = \psi_{m_d(v)} \circ f_1(z),$$

for all $z \in U'_{1,v}$.

Let $(\theta_v)_{v \in T(\lambda_0)}$ be such that, for $j = 1, 2$, the external marking Γ_j of the λ_0-renormalization of f_j is represented by $(\gamma_{j,v})_{v \in |T(\lambda_0)|}$ where $\gamma_{j,v} \subset R_{f_j}(\theta_v)$, for all $v \in |T(\lambda_0)|$. Then, we may adjust the collection $(\theta_v)_{v \in |T(\lambda_0)|}$ so that it extends to a larger one $(\theta_v)_{v \in |T^U(\lambda_0)|}$ such that $\theta_{m_d(v)} = d\theta_v$, for all $v \in |T^U(\lambda_0)|$. It follows that ψ_v maps the access to $K_v(f_1)$ determined by $R_{f_1}(\theta_v)$ onto the access to $K_v(f_2)$ determined by $R_{f_2}(\theta_v)$, for all $v \in |T^U(\lambda_0)|$. Hence, we can adjust the hybrid conjugacy ψ so that ψ_v maps $R_{f_1}(\theta_v) \cap U_{1,v}$ onto $R_{f_2}(\theta_v) \cap U_{2,v}$, for all v.

Let $\Lambda = (\Lambda_k)$ be a generator of λ_0 and $k_0 \geq 0$ be a separation depth. We may assume that every Julia critical element of λ_0 is a Λ_k-class for all $k \geq k_0$. As in the previous section, for each $v \in |T^U(\lambda_0)|$, let $L_k(v)$ be the combinatorial puzzle piece of depth $k \geq 0$ for Λ containing v. Denote by $P_k(f_j, v)$ the corresponding puzzle piece for (f_j, v). For each $v \in |T^U(\lambda_0)|$, take a neighborhood $V_{1,v}$ of $K_{f_1}(v) \cap \operatorname{int} P_{k_0}(f_1, v)$. and let $V_{2,v} = \psi_v(V_{1,v})$. We may choose $V_{1,v}$ sufficiently small so that

$$V_{j,v} \subset U_{j,v} \cap \operatorname{int} P_{k_0}(f_j, v).$$

Denote by $\operatorname{supp}(f_j, \Lambda_{k_0})$ the set formed by the landing points of the external rays of f_j with arguments in $\operatorname{supp}(\Lambda_{k_0})$. Shrinking $V_{j,v}$, if necessary, we may also assume that

$$\overline{V_{j,v}} \cap \partial P_{k_0}(f_j, v) \subset \operatorname{supp}(f_j, \Lambda_{k_0}).$$

In particular, for all $v \neq v' \in |T^U(\lambda_0)|$, the neighborhoods $V_{1,v}$ and $V_{j,v'}$ are disjoint and $\overline{V_{j,v}} \cap \overline{V_{j,v'}} \subset K_{f_j}(v) \cap K_{f_j}(v')$.

For $j = 1, 2$ let $\mathcal{V}_j = \bigcup_{v \in |T^U(\lambda_0)|} V_{j,v}$. Define $\Phi_0 : \mathcal{V}_1 \to \mathcal{V}_2$ by

$$\Phi_0 = \psi_v \text{ on } V_{1,v}.$$

In view of the above, we may extend Φ_0 quasiconformally to $\mathbb{C}$ such that:

- $\Phi_0(P_{k_0}(f_1, L)) = P_{k_0}(f_2, L)$ for all combinatorial pieces L of depth k_0.
- $\Phi_0(z) = \varphi_2^{-1} \circ \varphi_1(z)$ for all z in $\mathbb{C} \setminus D_{f_1}(r/d^{k_0})$ where φ_j denotes the Böttcher map of f_j.

- $\Phi_0(R_{f_1}(\theta_v)) = R_{f_2}(\theta_v)$ for all $v \in T^U(\lambda_0)$.

Note that Φ_0 is a conjugacy in an arcwise connected set S_0 which includes all the critical orbits of f_1. Also keep in mind that S_0 contains $\mathbb{C} \setminus D_{f_1}(r/d^{k_0})$ as well as $K_v(f_1)$ for all $v \in |T^U(\lambda_0)|$. Therefore, we may recursively define Φ_n, for $n \geq 1$, as the unique (quasiconformal) homeomorphism such that $\Phi_n \circ f_1 = f_2 \circ \Phi_{n-1}$ which agrees with Φ_{n-1} in the arcwise connected set $S_{n-1} = f_1^{-(n-1)}(S_0)$. It follows that the quasiconformal dilatation constant of Φ_n agrees with that of Φ_0 and, Φ_n is conformal in $\mathbb{C} \setminus D_{f_1}(r/d^{k_0+n})$ as well as in $f^{-n}(K_v(f_1))$, for all $v \in |T^U(\lambda_0)|$. Hence, after passing to a subsequence, Φ_n converges to a quasiconformal map Φ that conjugates f_1 and f_2 in the dense set $\bigcup S_n$, and therefore in $\mathbb{C}$. Moreover, Φ is conformal in

$$\bigcup_{n\geq 0} f^{-n}(\mathcal{K}(|T^U(\lambda_0)|) \cup (\mathbb{C} \setminus K(f_1)),$$

which is a full measure set in $\mathbb{C}$, in view of the Key Lemma. Therefore Φ is affine. In fact, Φ is the identity since it is a conjugacy between monic centered polynomials which is asymptotic to the identity at infinity. □

5.2. **Proof of the key lemma.** The rest of this section is devoted to prove Key Lemma 5.1. Let $\Lambda = (\Lambda_k)$ be a generator of λ_0 and let $k_0 \geq 0$ be a separation depth. Denote by $T^U(\lambda_0)$ the unreduced mapping schema of λ_0 as in the previous proof.

We introduce the following subsets of the Fatou critical orbit elements of λ_0 (i.e., $|T^U(\lambda_0)|$):

$$\begin{aligned} CO^0 &= \{v \in |T^U(\lambda_0)|;\ K_f(v) \cap \operatorname{supp}(f, \Lambda_{k_0+1}) = \emptyset\}, \\ CO^1 &= |T^U(\lambda_0)| \setminus CO^0. \end{aligned}$$

Denote by CO^{per} the set formed by the periodic Fatou critical elements of λ_0. Let

$$CO^{i,\mathrm{per}} = CO^i \cap CO^{\mathrm{per}},$$

for $i = 0, 1$ (see Figure 8).

In the following, for a set $X \subset \mathbb{C}$, we denote by $\operatorname{Comp}(X)$ denote the set of connected components of X and let $\mathcal{N}_\epsilon(X) = \{z \in \mathbb{C};\ d(z, X) < \epsilon\}$ be the ϵ-neighborhood of X.

Given a subset S of $|T^U(\lambda_0)|$, we let

$$\mathcal{K}_f(S) = \bigcup_{v\in S} K_f(v).$$

Fix $\epsilon_0 > 0$ sufficiently small, so that the neighborhoods $\mathcal{N}_{\epsilon_0}(\mathcal{K})$ are pairwise disjoint for the components $\mathcal{K}$ of $\mathcal{K}_f(CO^{\mathrm{per}})$. Recall that

$$F = K(f) \setminus \bigcup_{n\geq 0} \bigcup_{v\in|T(\lambda_0)|} f^{-n}(K_f(v)).$$

Lemma 5.4. *The following hold for almost every $z \in F$:*

(i) $\lim_{n\to\infty} \|(f^n)'(z)\| = \infty$ *with respect to the hyperbolic metric on* $\mathbb{C} \setminus PC(f)$;
(ii) $\lim_{n\to\infty} d(f^n(z), \mathcal{K}_f(CO^{1,\mathrm{per}})) = 0$;
(iii) *there exist $N = N(z) > 0$ and $\mathcal{K} \in \operatorname{Comp}(\mathcal{K}_f(CO^{1,\mathrm{per}}))$ such that $f^{N+n}(z) \in \mathcal{N}_{\epsilon_0}(f^n(\mathcal{K}))$. Moreover, if $f^{N+n}(z) \in \mathcal{N}_{\epsilon_0}(K_f(v))$ for $v \in CO^{1,\mathrm{per}}$, then $f^{N+n+1}(z) \in \mathcal{N}_{\epsilon_0}(K_f(m_d(v)))$.*

Proof. The assertion (i) is a stronger version of Mañé's Lemma by McMullen (see [Mc1, Theorem 3.6]).



FIGURE 8. Yoccoz puzzles of a separation depth. There are two periodic cycle of period two and three of Fatou components, which are colored black. The period two cycles corresponds to CO^0, and the period three cycles corresponds to CO^1.

In view of [Mc1, Theorem 3.9]), for almost every point $z \in J(f)$ we have that

$$\lim_{n\to\infty} d(f^n(z), PC(f)) = 0.$$

For all such $z \in F$,

$$\lim_{n\to\infty} d(f^n(z), \mathcal{K}_f(CO^{\mathrm{per}})) = 0$$

because $PC(f) \setminus \mathcal{K}_f(|T^U(\lambda_0)|)$ is a finite set consisting only of repelling periodic points and their backward images, and $f^n(\mathcal{K}_f(|T^U(\lambda_0))|) = \mathcal{K}_f(CO^{\mathrm{per}})$ for some $n \geq 0$. Since $\mathcal{K}_f(CO^{0,\mathrm{per}})$ and $\mathcal{K}_f(CO^{1,\mathrm{per}})$ are disjoint forward invariant compact sets, we have

$$\lim_{n\to\infty} d(f^n(z), \mathcal{K}_f(CO^{i,\mathrm{per}})) = 0$$

for some $i \in \{0, 1\}$.

Now assume

$$\lim_{n\to\infty} d(f^n(z), \mathcal{K}_f(CO^{0,\mathrm{per}})) = 0. \tag{2}$$

For each $v \in CO^0$, $K_f(v)$ is contained in the interior of the puzzle piece $P(L_{k_0+1}(v))$. Therefore we have $K_f(v) \in \mathrm{Comp}(\mathcal{K}_f(CO^{0,\mathrm{per}}))$ (i.e., $K_f(v)$ is a component of $\mathcal{K}_f(CO^{0,\mathrm{per}})$). By (2) and the continuity of f, it follows that there exist some $N \geq 0$ and $v \in CO^0$ such that for any $n \geq 0$, $f^{N+n}(z) \in P(L_{k_0+1}(m_d^n(v)))$, which is a neighborhood of $K_f(m_d^n(v))$. Let p be a period of v under iterations of m_d. Then $f^{pn}(f^N(z))$ belongs to a neighborhood of $K_f(v)$ for all $n \geq 0$. Since $K_f(v)$ is the filled Julia set of a polynomial-like map, it follows that $f^N(z) \in K_f(v)$, which contradicts $z \in F$. Therefore, we have proved (ii).

The assertion (iii) easily follows from (ii) and the continuity of f. □

Now we define, for each $v \in CO^{1,\mathrm{per}}$, an open set $\mathcal{N}'(v)$ (which is a slightly smaller set than $\mathcal{N}_{\epsilon_0}(K_f(v))$). Each $\mathcal{K} \in \mathrm{Comp}(\mathcal{K}_f(CO^{1,\mathrm{per}}))$ can be written as

$$\mathcal{K} = \bigcup_{m=0}^{M} K_f(v_m), \quad v_m \in CO^{1,\mathrm{per}}.$$

Then the set

$$I = \bigcup_{m \neq m'} (K_f(v_m) \cap K_f(v'_m)) \subset \mathrm{supp}(f, \Lambda_{k_0+1})$$

consists of repelling (pre)periodic points (see [Mc1, Theorem 7.3] and [In1, Proposition 3.4]). For each $x \in I$, take $\Theta(x) \subset \mathrm{Angle}(x) \cap \mathrm{supp}(\Lambda_{k_0})$ such that $d(\Theta(x)) = \Theta(f(x))$ and each component of the complement of

$$\Gamma(x) = \bigcup_{\theta \in \Theta(x)} R_f(\theta) \cup \{x\}$$

intersects exactly one of the $K_f(v_m)$ that contain x. Let $\Gamma = \bigcup_{x \in I} \Gamma(x)$ and let Γ_0 denote the union of the components of $\Gamma \cap \mathcal{N}_{\epsilon_0}(\mathcal{K})$ intersecting I. Then each component of $\mathcal{N}_{\epsilon_0}(\mathcal{K}) \setminus \Gamma_0$ intersects exactly one of $K_f(v_1), \ldots, K_f(v_M)$. Let us denote by $\mathcal{N}'(v_m)$ the component of $\mathcal{N}_{\epsilon_0}(\mathcal{K}) \setminus \Gamma_0$ containing $K_f(v_m) \setminus I$.

By construction,

$$\overline{\bigcup_{m=1}^{M} \mathcal{N}'(v_m)} = \overline{\mathcal{N}_{\epsilon_0}(\mathcal{K})} \tag{3}$$

and $\mathcal{N}'(v)$ ($v \in CO^{1,\mathrm{per}}$) are mutually disjoint.

Lemma 5.5. *Let $F_0 = \{z \in F;$ Lemma 5.4 holds$\}$. For $z \in F_0$, we have*

(i) *For all $n \geq N(z)$, there exists $v(z,n) \in CO^{1,\mathrm{per}}$ such that $f^n(z) \in \mathcal{N}'(v(z,n))$;*
(ii) *there exists arbitrarily large n such that $m_d(v(z,n)) \neq v(z,n+1)$;*
(iii) *there exists $\epsilon_1 > 0$ independent of z such that if $m_d(v(z,n)) \neq v(z,n+1)$, then $f^{n+1}(z) \in \mathcal{N}_{\epsilon_1}(I)$ and $f^n(z) \in \mathcal{N}_{\epsilon_1}((f^{-1}(I) \cap \mathcal{K}_f(CO^{1,\mathrm{per}})) \setminus I)$.*

Furthermore, ϵ_1 tends to zero as ϵ_0 tends to zero.

Proof. The assertion (i) follows from the equation (3) above. If (ii) does not hold, then we would have that $f^n(z) \in K_f(v(z,n))$, for sufficiently large n (since Λ is a generator for λ_0). This is a contradiction with $z \in F_0$. Finally, (iii) follows from the fact that $\mathcal{N}_{\epsilon_0}(K_f(v)) \setminus \mathcal{N}'(v)$ is contained in a small neighborhood of $I \cap K_f(v)$. □

Take a small neighborhood $\mathcal{O}$ of I such that $f(\mathcal{O}) \supset \overline{\mathcal{O}}$ and $f|_{\mathcal{O}}$ is injective. Since $CO(f) \setminus \mathcal{K}_f(CO^{\mathrm{per}})$ is finite, we may also assume the following:

(i) No critical value lies in $\mathcal{O}^* = \mathcal{O} \setminus I$,
(ii) If a component $\mathcal{O}_1$ of $f^{-1}(\mathcal{O})$ intersects $\mathcal{K}_f(CO^{1,\mathrm{per}})$, then $\mathcal{O}_1 \cap f^{-1}(I)$ and $\mathcal{O}_1 \cap CO(f)$ are contained in $\mathcal{K}_f(CO^{1,\mathrm{per}})$.
(iii) $\tilde{\mathcal{O}} = \mathcal{O}^*/f$ is a union of tori.

Figure 9 illustrates the statement of the following lemma.

Lemma 5.6. *There exist open subsets $\mathcal{U}_1$ and $\mathcal{U}_2$ of O such that the following statements hold:*

(i) $\mathcal{U}_2 \Subset \mathcal{U}_1 \Subset O$ *and* $\mathcal{U}_1$ *does not intersect* Γ.

(ii) *Each one of the open sets* $\mathcal{U}_1$ *and* $\mathcal{U}_2$ *is a disjoint union of finitely many topological disks.*

(iii) *Each component* A *of* $\mathcal{U}_1$ *contains exactly one component* B *of* $\mathcal{U}_2$ *and* $A \setminus \overline{B}$ *is an annulus.*

(iv) $\mathcal{U}_2/f \supset (O^* \cap K(f))/f$. *In other words, there exists* $\epsilon_2 > 0$ *such that for any* $z \in K(f) \cap \mathcal{N}_{\epsilon_2}(I)$, *there exists some* $N > 0$ *such that* $f^n(z) \in O$ *for any* n *with* $0 \leq n < N$, $f^N(z) \in \mathcal{U}_2$, *and the branch of* f^{-N} *sending* $f^N(z)$ *to* z *can be univalently defined on the component of* $\mathcal{U}_1$ *containing* $f^N(z)$

Proof. (See Figure 9.) Let $\tilde{K} = (K(f) \cap O^*)/f$ and $\tilde{\Gamma} = (\Gamma \cap O^*)/f$. Note that $\tilde{\Gamma}$ and $\tilde{K}$ are disjoint compact sets in $\tilde{O}$. Take a fundamental domain U for the covering projection $O \setminus \Gamma \to \tilde{O} \setminus \tilde{\Gamma}$. Let $\mathcal{U}_2 \Subset \mathcal{U}_1$ be neighborhoods of $U \cap K(f)$ in $O \setminus \Gamma$. It is easy to see that we can take $\mathcal{U}_1$ and $\mathcal{U}_2$ so that they satisfy the assertion. □

Proof of Key Lemma 5.1. It is enough to show that any $z \in F_0$ is not a Lebesgue density point of $K(f) \supset F_0$.

Let ϵ_1 and ϵ_2 be as in Lemma 5.5 and Lemma 5.6 respectively. We may assume $\epsilon_1 \leq \epsilon_2$. For $z \in F_0$, let

$$t(z) = \{n \geq N(z);\ m_d(v(z,n)) \neq v(z,n+1)\}.$$

By Lemma 5.5, we have $t(z)$ contains infinitely many elements. For $n \in t(z)$, let $O_1(z,n)$ be the component of $f^{-1}(O \setminus \Gamma)$ containing $f^n(z)$. By the condition (i) on O above, $f : O_1(z,n) \to O \setminus \Gamma$ is injective. (See Figure 9.)

For $z \in F_0$ and $n \in t(z)$, $f^{n+1}(z)$ lies in a ϵ_1-neighborhood of I by Lemma 5.5. Hence by Lemma 5.6, there exists some $n' > n$ such that $f^{n'}(z) \in \mathcal{U}_2$ and the branch of $f^{-n'+(n+1)}$ sending $f^{n'}(z)$ to $f^{n+1}(z)$ is univalently defined on $\mathcal{U}'_1(z,n)$, where $\mathcal{U}'_i(z,n)$ is the component of $\mathcal{U}_i$ containing $f^{n'}(z)$. Let $\tilde{\mathcal{U}}_i(z,n)$ be the component of $f^{-n'+n}(\mathcal{U}'_i(z,n))$ containing $f^n(z)$. Then $\tilde{\mathcal{U}}_i(z,n) \subset O_1(z,n)$ and $f^{n'+n} : \tilde{\mathcal{U}}_i(z,n) \to \mathcal{U}'_i(z,n)$ is a conformal isomorphism. Since $O_1(z,n) \cap PC(f) \subset O_1(z,n) \cap CO(f) = \emptyset$, there exists a univalent inverse branch $f^{-n} : \tilde{\mathcal{U}}_i(z,n) \to \mathcal{U}_i(z,n)$ sending $f^n(z)$ to z. Therefore, $f^n : \mathcal{U}_1(z,n) \to \tilde{\mathcal{U}}_1(z,n)$ and on O, $f^{n'} : \mathcal{U}_1(z,n) \to \mathcal{U}'_1(z,n)$ are conformal isomorphisms (condition (ii)).

Denote by $\iota_{z,n} : \mathcal{U}_1(z,n) \hookrightarrow \mathbb{C} \setminus PC(f)$ the inclusion and consider the following diagram:

$$\begin{tikzcd}
\mathcal{U}_1(z,n) \arrow[r, "f^n", "\cong"'] \arrow[d, hook, "\iota_{z,n}"] & \tilde{\mathcal{U}}_1(z,n) \arrow[r, "f^{n'-n}", "\cong"'] \arrow[d, hook] & \mathcal{U}'_1(z,n) \\
\mathbb{C} \setminus PC(f) \arrow[r, dotted, "f^n"] & \mathbb{C} \setminus PC(f).
\end{tikzcd}$$

Since $\lim_{n\to\infty} \|(f^n)'(z)\| = \infty$ with respect to the hyperbolic metric on $\mathbb{C} \setminus PC(f)$ and inclusion does not expand hyperbolic metric, we have $\lim_{n\to\infty} \|\iota'_{z,n}(z)\| = 0$ with respect to the corresponding hyperbolic metric. By Koebe distortion theorem, this implies that the diameter $\mathcal{U}_2(z,n)$ shrinks to zero with bounded distortion. Furthermore, since $f^{n'} : \mathcal{U}_1(z,n) \to \mathcal{U}'_1(z,n)$ is a conformal isomorphism sending $\mathcal{U}_2(z,n)$ to $\mathcal{U}'_2(z,n)$, by the Koebe distortion theorem applied to the inverse of the conformal isomorphism $f^{n'}|_{\mathcal{U}_1(z,n)}$), there exists some



FIGURE 9. Near I and its inverse image. If $m_d(v(z,n)) \neq v(z,n+1)$, $f^n(z)$ must be close to $f^{-1}(I) \setminus I$.

$C, C' > 0$ such that

$$\frac{\operatorname{Area}(\mathcal{U}_2(z,n) \setminus K(f))}{\operatorname{Area}(\mathcal{U}_2(z,n))} \geq C \frac{\operatorname{Area}(\mathcal{U}_2'(z,n) \setminus K(f))}{\operatorname{Area}(\mathcal{U}_2'(z,n))} \geq C'.$$

Here, we have only finitely many choices for $\mathcal{U}_i'(z,n)$ since each choice is a component of $\mathcal{U}_i$. This implies that C and C' are constants independent of $z \in F_0$ and $n \in t(z)$.

Therefore,

$$\lim_{n \in t(z) \to \infty} \frac{\operatorname{Area}(\mathcal{U}_2(z,n) \cap K(f))}{\operatorname{Area}(\mathcal{U}_2(z,n))} \leq 1 - C' < 1$$

and z is not a Lebesgue density point of $K(f)$. □

6. COMBINATORIAL TUNING

In this section we study the image of straightening and establish that, in a certain sense precised below, the straightening map is combinatorially onto.

Definition 6.1. A *rational lamination over a mapping schema* $T = (|T|, \sigma, \delta)$ is a collection of rational laminations $\lambda = (\lambda_v)_{v \in |T|}$ with $\delta(v)_* \lambda_v = \lambda_{\sigma(v)}$.

For $f \in \mathcal{C}(T)$, the *rational lamination* of f is the collection $\lambda_f = (\lambda_{f,v})_{v \in |T|}$ of equivalence relations in $\mathbb{Q}/\mathbb{Z}$ such that: θ and θ' are $\lambda_{f,v}$-equivalent if $R_f(v,\theta)$ and $R_f(v,\theta')$ land at the same point.

We can similarly define unlinked classes and the reduced mapping schema of a rational lamination over T. Hence we also define critical elements, post-critically finiteness and hyperbolicity of a rational lamination over T.

Definition 6.2. Let λ_0 and λ be invariant rational laminations. Assume λ_0 has non-empty reduced schema $T(\lambda_0)$ and $\lambda_0 \subset \lambda$. Let $\alpha = (\alpha_v : \overline{v} \to \mathbb{R}/\mathbb{Z})$ be an internal angle system for λ_0.

For each $v \in |T(\lambda_0)|$, let λ_v be the equivalence relation that identifies θ and θ' if and only if there exist $t \in \alpha_v^{-1}(\theta)$ and $s \in \alpha_v^{-1}(\theta')$ such that t and s are λ-equivalent. Then $\lambda' = (\lambda_v)_{v \in |T(\lambda_0)|}$ is a rational lamination over $T(\lambda_0)$. We say λ is the *combinatorial straightening* of λ with respect to (λ_0, α).

Theorem 6.3. *Consider an internally angled invariant rational lamination λ_0 with non-empty reduced schema and such that $\mathcal{R}(\lambda_0) \neq \emptyset$. Let $\chi : \mathcal{R}(\lambda_0) \to \mathcal{C}(T(\lambda_0))$ be the corresponding straightening map.*

If (λ_v) is a rational lamination over $T(\lambda_0)$, then there exists $f \in \mathcal{C}(\lambda_0)$ such that the combinatorial straightening of λ_f is equal to λ_v.

Moreover, we have the following:

- *If f above is λ_0-renormalizable, then the rational lamination of $\chi(f)$ is (λ_v).*
- *There exists an algebraic set $X = X(\lambda_0) \subset \operatorname{Poly}(d)$ of pure codimension one such that $\mathcal{R}(\lambda_0) \supset \mathcal{C}(\lambda_0) \setminus X$.*
- *There exists an algebraic set $Y = Y(\lambda_0) \subset \operatorname{Poly}(T(\lambda_0))$ of pure codimension one such that if $P \in \mathcal{C}(T(\lambda_0))$ is post-critically finite, then there exists $f \in \mathcal{R}(\lambda_0)$ such that $P = \chi(f)$ or P is a non-hyperbolic map contained in Y.*

Note that $\mathcal{C}(\lambda_0)$ might be contained in a proper algebraic set, so it can happen that $\mathcal{C}(\lambda_0)$ is contained in X and $\mathcal{R}(\lambda_0) = \emptyset$. Each irreducible component of the algebraic sets X and Y is defined by the existence of a parabolic periodic point of a given period, or a preperiodic critical point of given preperiod and eventual period.

The proof of the above theorem is at the end of this section, as well as that of Theorem C. As preliminary work for the proof we start by studying the action of straightening on rational laminations (see Lemma 6.4). Then we find an inverse for this action in Theorem 6.6. The last of the necessary ingredients is the characterization of laminations such that $\mathcal{R}(\lambda_0) \neq \emptyset$, contained in Proposition 6.7, which is based on the sufficient condition for a polynomial to belong to $\mathcal{R}(\lambda_0)$ given in Lemma 6.10.

6.1. **Combinatorial straightening and renormalized lamination.**

Lemma 6.4. *Assume that λ_0 is an invariant rational lamination with non-empty reduced schema. Let $\alpha = (\alpha_v : \overline{v} \to \mathbb{R}/\mathbb{Z})$ be an internal angle system for λ_0. Consider the corresponding straightening map $\chi : \mathcal{R}(\lambda_0) \to \mathcal{C}(T(\lambda_0))$. Given $f \in \mathcal{R}(\lambda_0)$ and let $\lambda'_f = (\lambda_v)_{v \in |T(\lambda_0)|}$ be the combinatorial straightening of λ_f.*

Then, for all $v \in |T(\lambda_0)|$,

$$\lambda_{\chi(f),v} = \lambda_v.$$

Proof. Let $g = (f^{\ell_v} : U'_v \to U_{\sigma_\lambda(v)})$ be a λ_0-renormalization. Let $P = \chi(g) : |T(\lambda_0)| \times \mathbb{C} \to |T(\lambda_0)| \times \mathbb{C}$ be the straightening of g via the fiberwise quasiconformal conjugacy $\psi_v : U_v \to \psi_v(U) \subset \{v\} \times \mathbb{C}$. Let $\varphi_v : \mathbb{C} \setminus K(P, v) \to \mathbb{C} \setminus \Delta$ be the Böttcher map. Given $v \in |T(\lambda_0)|$ and $\theta \in \overline{v} \cap \mathbb{Q}/\mathbb{Z}$, denote by $\beta_v(\theta) \in \mathbb{R}/\mathbb{Z} \cong \partial\Delta$ the "landing point" of the arc $\varphi_v \circ \psi_v(R_f(\theta) \cap U_v)$. That is, the closure of this arc intersects $\partial\Delta$ at a point with argument $\beta_v(\theta)$.

We claim that β_v and α_v coincide in $\overline{v} \cap \mathbb{Q}/\mathbb{Z}$. In fact, these maps at least coincide at one point, since ψ_v respects external markings. That is, if $\alpha_v(\theta_0) = 0$, then $\beta_v(\theta_0) = 0$. Now we observe that the set of arguments where α_0 and β_0 coincide is forward invariant. More precisely, let $A \subset \overline{v}$ be the largest set such that $m_d^{\ell_v}(A) = \{\theta\}$ where $\alpha_{\sigma_\lambda(v)}(\theta) = \beta_{\sigma_\lambda(v)}(\theta) = 0$. Notice that $\beta_{\sigma_\lambda(v)} \circ m_d = m_{\delta_\lambda(v)} \circ \beta_v$ and $\alpha_{\sigma_\lambda(v)} \circ m_d = m_{\delta_\lambda(v)} \circ \alpha_v$. Therefore, $\alpha_v(A) = \beta_v(A)$. Also, the cyclic order of the arguments in A is preserved by each of the maps α_v and β_v. Therefore, $\alpha_v(\theta) = \beta_v(\theta)$, for all $\theta \in A$. We may recursively repeat this argument in order to conclude that α_v and β_v coincide at a set C such that $\alpha_v(C)$ is dense. It follows that α_v and β_v agree on $\overline{v} \cap \mathbb{Q}/\mathbb{Z}$.

We conclude that $\lambda_v(P) \supset \lambda_v$, by Lindelöf Theorem.

Now consider two distinct λ_v-classes A_1 and A_2 and denote by z_1 and z_2 the landing points, in $K_f(v)$, of the rays with arguments in A_1 and A_2, respectively. Since ψ_v is injective, the landing points $\psi_v(z_1)$ and $\psi_v(z_2)$ of the external rays of P in the v-fiber, with arguments in $\alpha_v(A_1)$ and $\alpha_v(A_2)$ are distinct. Therefore, $\lambda_v(P) = \lambda_v$. □

6.2. **Combinatorial Tuning.**

In the classical context of quadratic polynomials, "tuning" is the inverse of straightening. Below we will show that, in a combinatorial sense, it is possible to find such an inverse.

Let us first introduce the pull-back of a relation. For a map $\varphi : A \to B$ and a relation λ on a set B, we say that $a_1 \in A$ is $\varphi^*\lambda$-related to $a_2 \in A$ if and only if $\varphi(a_1)$ is λ-related to $\varphi(a_2)$.

Definition 6.5. Let λ_0 be a d-invariant lamination with internal angle system $\alpha = (\alpha_v : \overline{v} \to \mathbb{R}/\mathbb{Z})$. Also, let $(\lambda_v)_{v \in |T(\lambda_0)|}$ be an invariant rational lamination over $T(\lambda_0)$.

Given an infinite λ_0-unlinked class v, denote by $n_v \geq 0$ the smallest integer such that $v' = m_d^{n_v}(v) \in |T(\lambda_0)|$. Let $\pi_v = \alpha_{v'} \circ m_d^{n_v} : \overline{v} \to \mathbb{R}/\mathbb{Z}$ and $\lambda_v^* = \pi_v^*(\lambda_{v'})$.

We define the *combinatorial tuning* $\mathcal{T}_\alpha(\lambda_0, (\lambda_v))$ as the smallest equivalence relation in $\mathbb{Q}/\mathbb{Z}$ containing:

(i) λ_0 and,
(ii) λ_v^*, for all infinite λ_0-unlinked classes v.

Theorem 6.6. *Let λ_0 be a d-invariant rational lamination, α be an internal angle system of λ_0 and $(\lambda_v)_{v\in|T(\lambda_0)|}$ be an invariant rational lamination over $T(\lambda_0)$. Then their combinatorial tuning $\lambda' = \mathcal{T}_\alpha(\lambda_0, (\lambda_v))$ is a d-invariant rational lamination containing λ_0. Moreover, the following statements hold:*

(i) *If λ_0 and (λ_v) are hyperbolic, then λ' is hyperbolic.*
(ii) *If λ_0 and (λ_v) are post-critically finite, then λ' is post-critically finite.*
(iii) *If $f \in \mathcal{R}(\lambda_0)$ has rational lamination λ', then the rational lamination of $\chi_{\lambda_0}(f)$ is (λ_v), where $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \to \mathcal{C}(T(\lambda_0))$ is the straightening map associated to the internal angle system α.*

Proof. Let $\mathcal{A}$ be the set of non-trivial λ_0-equivalence classes and let $\mathcal{B}$ be the union of the sets of non-trivial λ_v^*-equivalence classes for all infinite λ_0-unlinked classes v. A non-trivial λ'-equivalence class E is a maximal set of the form

$$E = \bigcup_{n=1}^{N} A_n \tag{4}$$

for $A_n \in \mathcal{A} \cup \mathcal{B}$ such that for any n, m, there exists a sequence $n_0 = n, n_1, \dots, n_k = m$ such that A_{n_j} and $A_{n_{j+1}}$ intersect for all $j = 1, \dots, k-1$. Finiteness of E follows from the fact the there exists ℓ and p such that $m_d^{\ell}(\theta) = m_d^{\ell+p}(\theta)$ for all $\theta \in E$, since the same holds for $E \in \mathcal{A} \cup \mathcal{B}$. Thus it is clear that every λ'-equivalence class is finite.

Observe that if $A, B \in \mathcal{A} \cup \mathcal{B}$ are disjoint, then A and B are unlinked. Consider $A_1, A_2, B_1, B_2 \in \mathcal{A} \cup \mathcal{B}$ such that A_1 intersects A_2 and B_1 intersects B_2, and $A_1 \cup A_2$ and $B_1 \cup B_2$ are disjoint. Then since A_j is unlinked with both B_1 and B_2, B_1 and B_2 are contained in the same component of $\mathbb{R}/\mathbb{Z} \setminus A_j$. Therefore, $A_1 \cup A_2$ and $B_1 \cup B_2$ are unlinked. We repeat this argument and conclude that λ'-equivalence classes are pairwise unlinked.

To see that λ' is closed, consider a sequence of pairs (θ_n, θ'_n) such that $\theta_n \sim_{\lambda'} \theta'_n$ and $\theta_n \to \theta$, $\theta'_n \to \theta'$ as $n \to \infty$. If $\theta_n \sim_{\lambda_0} \theta'_n$ for infinitely many n, then $\theta \sim_{\lambda_0} \theta'$ because λ_0 is closed, so it follows that $\theta \sim_{\lambda'} \theta'$. If there exists an infinite λ_0-unlinked class v such that $\theta_n, \theta'_n \in v$ for infinitely many n, then similarly we have $\theta \sim_{\lambda'} \theta'$ because λ_v^* is closed. Otherwise, by passing to a subsequence, we may assume that the convergences $\theta_n \nearrow \theta$ and $\theta'_n \searrow \theta'$ are monotone, that θ_n and θ'_n lie in an infinite λ_0-unlinked class v_n, and that v_n $(n = 1, 2, \dots)$ are pairwise disjoint. Then for each n, there exist $\iota_n \in (\theta_n, \theta_{n+1})$, $\iota'_n \in (\theta'_n, \theta'_{n+1})$ such that $\iota_n \sim_{\lambda_0} \iota'_n$ and that v_n and v_{n+1} lie in different components of $\mathbb{R}/\mathbb{Z} \setminus \{\iota_n, \iota'_n\}$. Therefore $\iota_n \nearrow \theta$ and $\iota'_n \searrow \theta'$, thus $\theta \sim_{\lambda_0} \theta'$.

To prove the d-invariance of λ', we start with the following observation: Let v be an infinite λ_0-unlinked class. Then $\pi_v : \overline{v} \to \mathbb{R}/\mathbb{Z}$ in Definition 6.5 induces a homeomorphism $\overline{v}/\lambda_0 \to \mathbb{R}/\mathbb{Z}$. Furthermore, $\pi_{m_d(v)} \circ m_d \circ \pi_v^{-1} : \mathbb{R}/\mathbb{Z} \to \mathbb{R}/\mathbb{Z}$ is well-defined equal to $m_{\delta(v)}$, and $\delta(v)_*\lambda_{v'} = \lambda_{w'}$, where $v' = m_d^{n_v}(v)$, $w = m_d(v)$ and $w' = m_d^{n_w}(w)$. Therefore, a λ_v^*-class (resp. λ_v^*-unlinked class) $A \subset \overline{v}$ is mapped by m_d to a $\lambda_{m_d(v)}^*$-class (resp. λ_v^*-unlinked class) and it is one-to-one if v is not critical, and $\delta(\alpha_v(A))$-to-one if v is critical.

Let E be a λ'-class of the form (4). Then its image

$$m_d(E) = \bigcup_{n=1}^{N} m_d(A_n)$$

is contained in some λ'-class E'. If $E' \setminus m_d(E)$ is nonempty, then there exist some $A' \in \mathcal{A} \cup \mathcal{B}$ intersecting both $E' \setminus m_d(E)$ and $m_d(E)$. If $A' \in \mathcal{A}$ then there exists a λ_0-class

A such that $m_d(A) = A'$ and A intersects E. Moreover, A is not contained in E because $m_d(A) = A' \not\subset m_d(E)$. This contradicts that E is maximal. Similarly, if A' is a $\lambda^*_{v'}$-class for an infinite λ_0-unlinked class v', then there exist an infinite λ_0-unlinked class v and a λ^*_v-class A such that $m_d(A) = A'$ and A intersects E but not contained in E, that is a contradiction. Therefore, $m_d(E) = E'$ is a λ'-equivalence class.

Now we show that $m_d : E \to m_d(E)$ is consecutive preserving. First, consider a λ'-unlinked class L. Then either L is a finite λ_0-unlinked class, or L is contained in an infinite λ_0-unlinked class. If L is a finite λ_0-unlinked class, then $m_d(L)$ is also a finite λ_0-unlinked class, which is again a λ'-unlinked class. On the contrary, if L is contained in an infinite λ_0-unlinked class v, then L is a λ^*_v-unlinked class. By construction, $m_d(L)$ is a $\lambda^*_{m_d(v)}$-unlinked class, which is also a λ'-unlinked class.

Let (θ_1, θ_2) be a component of $\mathbb{R}/\mathbb{Z} \setminus E$. Then there exist $\theta'_i \in (\theta_1, \theta_2)$ arbitrarily close to θ_i such that θ'_1 and θ'_2 are λ'-unlinked. In fact, if otherwise, there exists some $\varepsilon > 0$ such that any $\theta'_1 \in (\theta_1, \theta_1+\varepsilon)$ and $\theta'_2 \in (\theta_2-\varepsilon, \theta_2)$ are not λ'-unlinked. Hence there exist sequence $t_n \in (\theta_1, \theta_1 + 1/n)$ and $s_n \in (\theta_1 + \varepsilon, \theta_2 - \varepsilon)$ such that t_n and s_n are λ'-equivalent. By passing to a subsequence, we may assume s_n converges to some $s \in [\theta_1+\varepsilon, \theta_2-\varepsilon] \subset (\theta_1, \theta_2)$. Since we already proved that λ' is closed, it follows that θ_1 and s are λ'-equivalent and $s \in E$, that is a contradiction.

Thus for any $\theta \in (d\theta_1, d\theta_2)$, there exists some $\theta'_1, \theta'_2 \in (\theta_1, \theta_2)$ such that θ'_1 and θ'_2 are λ'-unlinked and $\{\theta'_1, \theta'_2\}$ and $\{\theta', \theta_1\}$ are linked (not unlinked) for any $\theta' \in m_d^{-1}(\theta) \cap (\theta_1, \theta_2)$. Since $d\theta'_1$ and $d\theta'_2$ are also λ'-unlinked, θ is not λ'-equivalent to $d\theta_1$. Therefore $(d\theta_1, d\theta_2)$ is a component of $\mathbb{R}/\mathbb{Z} \setminus m_d(E)$ and we have proved $m_d : E \to m_d(E)$ is consecutive preserving.

It follows that, λ' is d-invariant.

For (i), let us further assume that λ_0 and (λ_v) are post-critically finite. Let v' be a finite λ'-unlinked class. Then v' is either a finite λ_0-unlinked class, or a finite λ^*_v-unlinked class for some infinite λ_0-unlinked class v. Hence v' is not critical by assumption. Therefore, λ' is post-critically finite.

For (ii), if λ_0 and (λ_v) are hyperbolic, then for each Fatou critical element $\tilde{v}'$ of λ_v, $v' = \alpha_v^{-1}(\tilde{v}')$ is a critical element of λ' with $\delta(v') = \delta(\tilde{v}')$. Also, since

$$\sum_{\tilde{v}' \in \mathrm{Crit}((\lambda_v))} (\delta(\tilde{v}') - 1) = d - 1,$$

all the critical elements of λ' are infinite λ' unlinked classes. Therefore, λ' is also hyperbolic.

For (iii) assume that $f \in \mathcal{R}(\lambda_0)$ is such that $\lambda_f = \lambda'$. Then $\alpha_v^* \lambda_{\chi(f),v} = \lambda'|_{\overline{v}}$ for $v \in |T(\lambda_0)|$ by Lemma 6.4, hence it suffices to show that $\lambda'|_{\overline{v}} = \lambda^*_v$. Since we have $\lambda^*_v \subset \lambda'|_{\overline{v}}$ by definition, we need only show the following: Let E be a λ'-class of the form (4) and if A_1 intersects v, then $A_2, \dots, A_N$ do not intersect v.

For $n \geq 2$, take a sequence $n_0 = 1, n_1, \dots, n_K = n$ such that A_{n_k} intersects $A_{n_{k+1}}$ for $k = 0, \dots, K-1$. We may assume $A_{n_k} \neq A_{n_{k'}}$ if $k \neq k'$. Furthermore, we may assume

$$\#(A_{n_k} \cap A_{n_{k+1}}) = 2 \text{ for any } k. \tag{5}$$

In fact, at least one of A_{n_k} and $A_{n_{k+1}}$, say A_{n_k}, is a $\lambda^*_{v'}$-class for some v'. Then $A_{n_{k+1}}$ is not a $\lambda^*_{v'}$-class, so it is a λ-class or a $\lambda^*_{v''}$-class for some v'' whose boundary $\partial v''$ intersects $\partial v'$. If $A_{n_{k+1}}$ is a λ-class, then $A_{n_{k+1}} \cap \partial v$ contains exactly two points and $A_{n_{k+1}}$ is contained in A_{n_k}. If $A_{n_{k+1}}$ is a $\lambda^*_{v''}$-class, then $\partial v' \cap \partial v''$ is contained in a λ-class A, so we may add A in the sequence between A_{n_k} and $A_{n_{k+1}}$ to satisfy the condition (5).

Since A_{n_0} is a λ_v^*-class, A_{n_1} is not. In particular, A_{n_1} is contained in a component of $\mathbb{R}/\mathbb{Z} \setminus v$.

For $k \geq 1$, assume A_{n_k} is contained in a component of $\mathbb{R}/\mathbb{Z} \setminus v$ and $A_{n_{k-1}}$ is contained in a component of $\mathbb{R}/\mathbb{Z} \setminus A_{n_k}$ containing v. Then $A_{n_{k+1}}$ lies in a component of $\mathbb{R}/\mathbb{Z} \setminus A_{n_k}$ different from the component containing $A_{n_{k-1}}$. Since otherwise we have $\#(A_{n_{k-1}} \cap A_{n_k} \cap A_{n_{k+1}}) \geq 2$, which implies that two of them are equal. In particular, A_{n_k} is contained in a component of $\mathbb{R}/\mathbb{Z} \setminus A_{n_{k+1}}$ containing $A_{n_{k-1}}$ and v. Therefore, by induction, $A_n = A_{n_K}$ is contained in a component of $\mathbb{R}/\mathbb{Z} \setminus v$ which contains A_{n_1}, that is, $A_n \cap v = \emptyset$. Therefore, the rational lamination of $\chi_{\lambda_0}(f)$ is (λ_v).

□

6.3. **Renormalizable set.** Our aim now is to characterize laminations with non-empty renormalizable set.

Proposition 6.7. *Consider a d-invariant rational lamination such that* $T(\lambda_0) \neq \emptyset$. *Then the following statements are equivalent:*

(i) $\mathcal{R}(\lambda_0) \neq \emptyset$.

(ii) *If A is a critical* λ_0*-class and L is an infinite unlinked class in the forward orbit of a critical element, then* $A \cap \overline{L} = \emptyset$.

Before proving the above result let us introduce an important class of rational laminations.

Definition 6.8 (Primitive invariant laminations)**.** We say a d-invariant rational lamination is *primitive* if there do not exist infinite λ-unlinked classes L and L' and λ-class A such that $L \neq L'$ and both $\overline{L} \cap A$ and $\overline{L'} \cap A$ are nonempty.

As a direct consequence of the proposition we obtain the following result.

Corollary 6.9. *If* λ_0 *is a primitive rational lamination and* $|T(\lambda_0)|$, *then* $\mathcal{R}(\lambda_0) \neq \emptyset$.

To prove the proposition, we start with the lemma below, which gives sufficient conditions for a polynomial $f \in C(\lambda_0)$ to be in $\mathcal{R}(\lambda_0)$. Then the reader may find the proof of the proposition after the proof of the lemma.

Lemma 6.10. *Consider an invariant rational lamination* λ_0 *such that the following holds: if A is a critical* λ_0*-class and L is an infinite unlinked class in the forward orbit of a critical element, then* $A \cap \overline{L} = \emptyset$. *Let* $\Lambda = (\Lambda_k)$ *be a combinatorial Yoccoz puzzle which is a generator of* λ_0.

There exists $K = K(\lambda_0) > 0$ *such that if* $f \in C(\lambda_0)$ *is such that,*

- *for all* $\theta \in \operatorname{supp} \Lambda_0$, *the external ray* $R_f(\theta)$ *lands at a repelling periodic point, and*
- *for all* $\theta \in \operatorname{supp} \lambda_K$, *the landing point of the external ray* $R_f(\theta)$ *is not a critical point,*

then $f \in \mathcal{R}(\lambda_0)$.

By this lemma, we conclude the existence of X in Theorem 6.3:

Corollary 6.11. *Let* λ_0 *be an invariant rational lamination. Then there exists an algebraic set* $X \subset \operatorname{Poly}(d)$ *of pure codimension one such that* $\mathcal{R}(\lambda_0) \supset C(\lambda_0) \setminus X$.

Proof. This is a direct consequence of Lemma 6.10 if λ_0 satisfies the assumption.

Otherwise $C(\lambda_0)$ is contained in a proper algebraic subset X of Poly(d) because every $f \in C(\lambda_0)$ has a preperiodic critical point of preperiod and eventual period depending only on λ_0. (Indeed we have $\mathcal{R}(\lambda_0) = C(\lambda_0) \setminus X = \emptyset$ by Proposition 6.7.) □

Proof of Lemma 6.10. We claim that under the hypothesis of the lemma $K_f(v) = \bigcap P_k(f, v)$. Directly from the definitions it follows that $K_f(v) \subset \bigcap P_k(f, v)$. Now if $z \notin K_f(v)$, then there exists a sector S bounded by λ_0-equivalent rays with arguments in $\overline{v}$ that is disjoint from $K_f(v)$ and such that $z \in S$. Since Λ is a generator, there exists k and a sector S' bounded by Λ_k-equivalent arguments such that $z \in S' \subset S$ by Proposition 4.9. Thus, $z \notin P_k(f, v)$.

Now in order to show that $f \in \mathcal{R}(\lambda_0)$ we apply the thickening procedure (cf. [Mi2]). More precisely, let k_0 be a separation depth. By Proposition 4.9 and the assumption, we can take k_0 sufficiently large so that for all $k > k_0$ and for all v in the forward element of a Fatou critical element of λ_0, there exists no critical λ_k-class A which intersects $\overline{L_k(v)}$, where $L_k(v) \in \mathcal{L}_k(\Lambda)$ is the combinatorial puzzle piece of depth k containing v. Then for all v in the forward element of a Fatou critical element of λ_0, and all $k > k_0$, if the landing point of $R_f(\theta)$ for every $\theta \in \partial L_k(v)$ is neither parabolic nor critical, then the thickening procedure gives Jordan domains $\hat{P}_k(f, v)$ such that:

(i) $P_k(f, v) \subset \hat{P}_k(f, v) \Subset \hat{P}_{k-1}(f, v)$,
(ii) $f : \hat{P}_k(f, v) \to \hat{P}_{k-1}(f, \sigma_\lambda(v))$ is a proper map of degree $\delta_\lambda(v)$ (compare Proposition 3.7 (iv)(e)) and,
(iii) $\bigcap \hat{P}_k(f, v) = \bigcap P_k(f, v)$.

It follows that if we take $U'_v = \hat{P}_{k_0+\ell_v}(f, v)$ and $U_v = \hat{P}_{k_0}(f, v)$ for all $v \in |T(\lambda_0)|$, then $g = (f^{\ell_v} : U'_v \to U_{\sigma_\lambda(v)})$ is the desired polynomial-like map over $T(\lambda_0)$ with the appropriate fiberwise connected Julia set. Therefore if $f \in C(\lambda_0)$ satisfies the assumption for

$$K = k_0 + \max_{v \in |T(\lambda_0)|} \ell_v,$$

then $f \in \mathcal{R}(\lambda_0)$. □

Proof of Proposition 6.7. (i) $\Longrightarrow$ (ii). Assume that $f \in \mathcal{R}(\lambda_0)$. Then there exists a polynomial-like map g over $T(\lambda_0)$ such that $K(g, v) = K_f(v)$ for all $v \in |T(\lambda_0)|$. For each $v \in |T(\lambda_0)|$, there exist a neighborhoods $U'_v \Subset U_v$ such that $f^{\ell_v} : U'_v \to U_{\sigma_{\lambda_0}(v)}$ is a proper map of degree $\delta_\lambda(v)$. In order to conclude that (ii) holds, we proceed by contradiction and suppose that there exists a critical λ_0-class A such that $A \cap m_d^\ell(\overline{v}) \neq \emptyset$ for some $0 \leq \ell < \ell_v$. Note that there are exactly $\delta_{\lambda_0}(v)$ λ_0-classes $B_1, \dots, B_{\delta_{\lambda_0}(v)}$ intersecting $\overline{v}$ that map into the λ_0-class $m_d^{\ell_v - \ell}(A)$ under $m_d^{\ell_v}$. Let $A_1, \dots, A_{\delta_\lambda(v)}$ be the corresponding λ_f-classes and $t \in m_d^{\ell_v - \ell}(A)$. It follows that the set

$$\{t' \in A_1 \cup \dots \cup A_{\delta_{\lambda_0}(v)}; m_d^{\ell_v}(t') = t\}$$

has cardinality strictly greater than $\delta_{\lambda_0}(v)$, since $m_d^\ell(A_j) = A$ for at least one j. Therefore, the neighborhood U'_v of $K_f(v)$ contains the portion inside $D_f(r)$, for r sufficiently small, of at least $\delta_{\lambda_0}(v) + 1$ rays which map onto the same ray under f^{ℓ_v}. Hence $f^{\ell_v} : U'_v \to U_{\sigma_{\lambda_0}(v)}$ is not a degree $\delta_{\lambda_0}(v)$ map.

(ii) $\Longrightarrow$ (i). Assume that (ii) holds, according to Lemma 7.2, Corollary 7.3 and the proof of Theorem 1.1 in [Ki], there exists a polynomial $f \in C(d)$ with rational lamination $\lambda_f = \lambda_0$ and no neutral cycles. Hence, we may apply the previous lemma to a generator of λ_0 given by Lemma 4.10. □

6.4. **Proof of Theorem 6.3.** Denote by α the internal angle system. By Theorem 6.6, and Lemma 7.2, Corollary 7.3 and the proof of Theorem 1.1 in [Ki], there exists $f \in C(\lambda_0)$ with no neutral cycles and rational lamination $\lambda_f = \mathcal{T}_\alpha(\lambda_0, (\lambda_v))$ such that every Fatou critical point has finite forward orbit. By Proposition 6.7 and Lemma 6.10, it follows that there exists an algebraic set $Y \subset \mathrm{Poly}(T(\lambda_0))$ of pure codimension one such that $f \in \mathcal{R}(\lambda_0)$, except when $P \in Y$. More precisely, the assumption of Lemma 6.10 does not hold if and only if the landing point of $R_P(v, \alpha_v(\theta))$ is a critical point for some $v \in CO(\lambda_0)$ and $\theta \in \operatorname{supp} \Lambda_K \cap \partial v$, where K is in Lemma 6.10. In particular, P has a critical point in the Julia set, i.e., P is not hyperbolic. Except for that case, from Theorem 6.6, we conclude that $\lambda_{\chi(f),v} = \lambda_v$ for all $v \in |T(\lambda_0)|$.

Now if $P \in \mathcal{C}(T(\lambda_0))$ is post-critically finite, then the standard pull-back argument shows that P is uniquely determined by its rational lamination. Taking f as in the previous paragraph, such that $\lambda_{\chi(f)} = \lambda_P$, then we have that $\chi(f) = P$. □

6.5. **Proof of Theorem C.** Let λ_0 be a post-critically finite rational lamination. Consider a component $\mathcal{H}$ of $\mathrm{Hyp}\,\mathcal{C}(T(\lambda_0))$. Our aim now is to show that the image of χ_{λ_0} contains $\mathcal{H}$.

Consider the unique post-critically finite polynomial map $P_0 \in \mathcal{H}$ [Mi4, Corollary 4.2]. Since P_0 is hyperbolic, by Theorem 6.3, there exists $f_0 \in \mathcal{R}(\lambda_0)$ such that $\chi_{\lambda_0}(f_0) = P_0$. Note that $\mathcal{R}(\lambda_0) \subset \mathcal{C}(\lambda_0)$ is contained in an algebraic set Y of $\mathrm{Poly}(d)$. Moreover, the dynamics of f_0 over its Julia set is structurally stable in Y. Call $\mathcal{H}'$ the component of structurally stable maps in Y containing f_0. We claim that $\chi_{\lambda_0}(\mathcal{H}') = \mathcal{H}$.

In order to prove that $\chi_{\lambda_0}(\mathcal{H}') = \mathcal{H}$ we will employ Milnor's parametrization of $\mathcal{H}$. Let $T_0 = (|T_0|, \sigma_0, \delta_0)$ be the reduced mapping schema of P_0. Namely, $|T_0|$ is the set formed by the critical points of P_0, $\sigma_0 : |T_0| \to |T_0|$ is the first return map under iterations of P_0, and $\delta_0 : |T_0| \to \mathbb{N}$ is the local degree of P_0 at its critical points. Consider the space $B(T_0)$ of all proper holomorphic maps

$$\beta : |T_0| \times \Delta \to |T_0| \times \Delta$$

of the form $\beta(v,z) = (\sigma_0(v), \beta_v(z))$ where $\beta_v : \Delta \to \Delta$ is of degree $\delta_0(v)$ and the following hold:

- β_v is *boundary-rooted*, i.e., the extension of β_v to $S^1 = \partial\Delta$ satisfies $\beta_v(1) = 1$;
- if v is periodic under σ_0, then β_v is *fixed point centered*, i.e., $\beta_v(0) = 0$;
- if v is not periodic, then β_v is *critically centered*, i.e., the sum of its $\delta(v) - 1$ critical points (counted with multiplicity) is equal to zero.

According to Milnor [Mi4, Lemma 3.5] $B(T_0)$ is a topological cell. Now a map $\Phi : \mathcal{H} \to B(T_0)$ is introduced as follows. Given $P \in \mathcal{H}$, appropriate choices for the Riemann mappings, of the critical bounded Fatou components of P, "conjugate" the action of the first return map to that of an element $\Phi(P)$ of $B(T_0)$. Milnor showed that $\Phi : \mathcal{H} \to B(T_0)$ is a diffeomorphism [Mi4, Theorem 4.1']. Using the fact that hybrid conjugacies between the λ_0-renormalization of $f \in \mathcal{H}'$ and the polynomial maps $\chi(f) \in \mathcal{H}$ restrict to conformal isomorphisms of bounded Fatou components, the standard quasiconformal surgery argument shows that $\Phi \circ \chi : \mathcal{H}' \to B(T_0)$ is onto. Thus, $\chi : \mathcal{H}' \to \mathcal{H}$ is also surjective and, therefore, a bijection.

Since we have already shown that $\chi_{\lambda_0} : \mathcal{H}' \to \mathcal{H}$ is a continuous bijection, along the same lines of the proof of [Mi4, Theorem 5.1], it follows that $\chi_{\lambda_0} : \mathcal{H}' \to \mathcal{H}$ is in fact biholomorphic. □

7. COMPACTNESS

The main result of this section is to establish the following compactness result which generalizes [In3, Theorem 1.3].

Theorem 7.1. *If λ_0 is a primitive invariant rational lamination, then $\mathcal{C}(\lambda_0) = \mathcal{R}(\lambda_0)$ and this set is compact.*

The reader may find the proofs of Theorem 7.1 and Theorem D after the statements and proofs of the following two lemmas.

Primitive rational laminations are rich in non-trivial classes:

Lemma 7.2. *Let λ_0 be a non-trivial invariant rational lamination such that the support of λ_0 is contained in the m_d-grand orbit of a finite set. Then λ_0 is not primitive.*

Proof. By passing to an iterate, we may assume that the support of λ_0 is contained in the grand orbit of the finite set $F = \{j/(d-1); j = 0, \dots, d-2\}$. That is, $t \in F$ if and only if $m_d(t) = t$. We proceed by contradiction and assume that λ_0 is primitive.

Let $A_1, \dots, A_n$ be a complete list (without repetitions) of the λ_0-classes contained in F. Let f be a polynomial without neutral fixed points such that $\lambda_f = \lambda_0$. Let m be the number of non-trivial classes in F and observe that $n + m \leq d - 1$.

The union of the closure of the external rays with arguments in F cuts the complex plane into $d - n$ regions $U_1, \dots, U_{d-n}$, since

$$d - n = 1 + \sum_{j=1}^{n} (|A_j| - 1).$$

According to Goldberg and Milnor [GM], for all j, there exists exactly one fixed point z_j in the region U_j. These fixed points are not the landing point of rational rays since the lamination is supported in the grand orbit of F. Therefore, z_j is an attracting fixed point, for all j. For each $j = 1, \dots, d-n$, call L_j the fixed infinite λ_0-unlinked class such that $z_j \in K_f(L_j)$. By Proposition 3.7 (iv)(c), it follows that the boundary of L_j intersects a non-trivial class contained in F, say $A(L_j)$. Hence, for some $j \neq k$, we have that $A(L_j) = A(L_k)$. For otherwise, $d - n \leq m$, but $n + m \leq d - 1$. Therefore, λ_0 is not primitive which gives us the desired contradiction. □

Lemma 7.3. *Consider a non-trivial primitive d-invariant rational lamination λ_0. Let E be the m_d-grand orbit of a finite set and $\lambda = \lambda_0 \cap ((\mathbb{Q}/\mathbb{Z} \setminus E) \times (\mathbb{Q}/\mathbb{Z} \setminus E))$. Then λ_0 is the smallest closed equivalence relation in $\mathbb{Q}/\mathbb{Z}$ containing λ.*

Proof. Let λ' be the smallest closed equivalence relation in $\mathbb{Q}/\mathbb{Z}$ containing λ. It follows that λ' is a d-invariant rational lamination and $\lambda' \subset \lambda_0$.

Let L_0 be an infinite periodic λ'-unlinked class, say of period p. We claim that L_0 is a λ_0-unlinked class. For this, consider $\pi : \overline{L_0} \to \mathbb{R}/\mathbb{Z}$ such that $\pi \circ m_d^p = m_{d'} \circ \pi$ for some $d' \geq 2$. Let $\pi_* \lambda_0$ be the equivalence relation in $\mathbb{Q}/\mathbb{Z}$ that identifies θ and θ' if and only if there exists $t \in \pi^{-1}(\theta)$ and $s \in \pi^{-1}(\theta')$ such that s and t are λ_0-equivalent arguments. Observe that $\pi_* \lambda_0$ is a d'-invariant rational lamination with support contained in $\pi(E \cap \overline{L_0})$. By the previous lemma, $\pi_* \lambda_0$ is trivial or it is not primitive. The latter alternative is impossible, since taking the preimage of appropriate $\pi_* \lambda_0$-unlinked classes we would have that λ_0 is not primitive. Therefore, $\pi_* \lambda_0$ is trivial. Hence, L_0 is a λ_0-unlinked class.

Now let L_0 be a strictly preperiodic infinite λ'-unlinked class. We also claim that L_0 is a λ_0-class. In fact, let $\ell \geq 1$ be such that $m_d^\ell(L_0)$ is a periodic λ_0-unlinked class. If $\pi : \overline{L_0} \to \mathbb{R}/\mathbb{Z}$ is such that $\pi \circ m_d^\ell = m_{d'} \circ \pi$ for some $d' \geq 1$, then $m_{d'*}(\pi_* \lambda_0)$ is the trivial rational lamination. In particular, every non-trivial $\pi_* \lambda_0$-class is mapped, under $m_{d'}$ onto a singleton. Since these classes are unlinked, there are at most $d' - 1$ non-trivial $\pi_* \lambda_0$-classes. Therefore, either $\pi_* \lambda_0$ is trivial and L_0 is a λ_0-unlinked class, or λ_0 is not primitive. Thus, L_0 is a λ_0-unlinked class.

Now to prove that $\lambda' = \lambda_0$ just observe that otherwise we would have an infinite λ'-unlinked class that is not a λ_0-unlinked class. □

With the previous lemma we may now establish our compactness result.

Proof of Theorem 7.1. Assume that $f \in \mathcal{C}(\lambda_0)$. In order to show that $\mathcal{C}(\lambda_0) = \mathcal{R}(\lambda_0)$ we must prove that f is λ_0-renormalizable (i.e., $f \in \mathcal{R}(\lambda_0)$). More precisely, we must extract a polynomial-like map g over $T(\lambda_0)$ (see Definition 3.12). To extract a polynomial-like map from f we will apply the thickening procedure (Lemma 6.10) to an appropriate Yoccoz puzzle.

We start by finding an appropriate combinatorial Yoccoz puzzle. Let $F \subset \mathbb{Q}/\mathbb{Z}$ be the set formed by the arguments of external rays of f landing at parabolic or critical points. Let m be the minimum common multiple of the periods, under m_d, of the arguments in the forward orbit of F. Denote by $F' \subset \mathbb{Q}/\mathbb{Z}$ the set of all m_d-periodic arguments of period dividing m and, by E' the grand orbit of F'. From the previous lemma, there exist λ_0-classes $A_1, \dots, A_j$ contained in $\mathbb{Q}/\mathbb{Z} \setminus E'$ that separate the critical elements of λ_0. The support of the depth 0 puzzle will be $E_0 \subset \mathbb{Q}/\mathbb{Z}$, the set formed by all periodic arguments in the forward orbit of $A_1 \cup \dots \cup A_j$. More precisely, let λ be the restriction of λ_0 to $E_0 \times E_0$. For all $\ell \geq 1$, let $E_\ell = m_d^{-\ell}(E_0)$ and define Λ_ℓ as the restriction of λ_0 to $E_\ell \times E_\ell$. It follows that $\Lambda = (\Lambda_\ell)$ is a combinatorial Yoccoz puzzle admissible for λ_0. Note that, for some $k \geq 0$, we have that $E_k \supset A_1 \cup \dots \cup A_j$. In particular, Λ is a generator and such a number k is a separation depth for λ_0.

Since $E_\ell \cap E' = \emptyset$, for all $\theta \in E_\ell$, the landing points of the external ray of angle θ is a non-critical eventually repelling periodic point. By Lemma 6.10, we have that $f \in \mathcal{R}(\lambda_0)$.

Now we show that $\mathcal{C}(\lambda_0)$ is compact. Due to the compactness of $\mathcal{C}(d)$, we only have to show that $\mathcal{C}(\lambda_0)$ is closed in $\mathcal{C}(d)$. For this consider a sequence (f_n) of polynomials in $\mathcal{C}(\lambda_0)$ which converges to $f \in \mathcal{C}(d)$. Let $F \subset \mathbb{Q}/\mathbb{Z}$ be the arguments of the rays landing at parabolic periodic points of f or at critical points of f. Denote by E the m_d-grand orbit of F.

We claim that

$$\lambda_0 \cap ((\mathbb{Q}/\mathbb{Z} \setminus E) \times (\mathbb{Q}/\mathbb{Z} \setminus E)) \subset \lambda_f \cap ((\mathbb{Q}/\mathbb{Z} \setminus E) \times (\mathbb{Q}/\mathbb{Z} \setminus E)).$$

Assume that two arguments θ and θ' in $\mathbb{Q}/\mathbb{Z} \setminus E$ are λ_0-equivalent. Since the landing points of the external rays with arguments θ and θ' are repelling for f, they depend continuously on the polynomial in a neighborhood of f. Hence for n sufficiently large, θ and θ' are λ_{f_n}-equivalent if and only if they are λ_f-equivalent. Since $\lambda_{f_n} \supset \lambda_0$, we have proved that θ and θ' are λ_f-equivalent.

From the previous lemma and the fact that λ_f is closed in $\mathbb{Q}/\mathbb{Z} \times \mathbb{Q}/\mathbb{Z}$ we conclude that $\lambda_f \supset \lambda_0$. Hence, $f \in \mathcal{C}(\lambda_0)$. ◻

Proof of Theorem D. Assume that λ_0 is post-critically finite but not primitive. From the previously proven theorem, it is sufficient to show that: $\mathcal{R}(\lambda_0)$ is not compact and $\mathcal{C}(\lambda_0) \neq \mathcal{R}(\lambda_0)$.

Let $f_0 \in \mathcal{R}(\lambda_0)$ be hyperbolic with $\lambda_{f_0} = \lambda_0$. Since λ_0 is not primitive, there exist two periodic Fatou components Ω_1, Ω_2 such that $\partial\Omega_1 \cap \partial\Omega_2 = \{x_0\}$ where x_0 is a repelling periodic point of period dividing one of those of these two periodic Fatou components, say Ω_1. We may assume that f_0 has an attracting (not superattracting) periodic point y_0 in Ω_1. By Haïssinsky's pinching theorem [Ha1] [Ha4], there exist a continuous path of quasiconformal deformations of polynomials $(f_t)_{t \in [0,1)}$ such that

(i) f_t converges uniformly to a polynomial f_1, shrinking progressively some path γ connecting x_0 and y_0. There exists a semiconjugacy $\varphi : \mathbb{C} \to \mathbb{C}$ from f_0 to f_1 such that φ is a homeomorphism outside γ and its preimages, and φ sends γ to a point, as well as each component of the preimages of γ.

(ii) The point $x_1 = \varphi(\gamma)$ is a parabolic periodic point f_1 and its immediate basin is equal to $\varphi(\Omega_1 \setminus \Gamma)$ where Γ is the union of γ and all the preimages.

In particular, f_0 and f_1 are topologically conjugate on their Julia sets, by φ. Therefore, $f_t \in \mathcal{R}(\lambda_0)$ for $t \in [0,1)$ but $f_1 \in \mathcal{C}(\lambda_0)$ and $f_1 \notin \mathcal{R}(\lambda_0)$. Otherwise, a λ_0-renormalization g of f_1 would satisfy $\{x_1\} = K(g, v_1) \cap K(g, v_2)$ where v_i is the infinite λ_0-unlinked class such that $K(f_0, v_i) = \overline{\Omega_i}$. Therefore, x_1 would be repelling ([Mc1, Theorem 7.3] and [In1, Proposition 3.4]) which is a contradiction with (ii). □

Remark 7.4. For postcritically finite invariant rational laminations λ_0, the remaining problem is to characterize when $\mathcal{C}(\lambda_0)$ is compact. Note that $\mathcal{C}(\lambda_0)$ is known to be compact in some cases and non-compact in some other cases. For example, if λ_0 is of degree two, then $\mathcal{C}(\lambda_0)$ is always compact (it is a baby Mandelbrot set).

On the other hand, the following example shows that $\mathcal{C}(\lambda_0)$ is not compact in general.

Example 7.5. Let λ_0 be a rational lamination of $f_0(z) = z^3 + 3z/2$. The critical points $\pm i/\sqrt{2}$ of f_0 are fixed and the origin lies in the boundaries of their immediate basin of attraction. Since f_0 is real and $K(f_0) \cap \mathbb{R} = \{0\}$, the external rays of angles 0 and 1/2 are the positive and negative real line respectively, so 0 and $\frac{1}{2}$ are λ_0-equivalent.

For $\mu = \zeta + i\xi \neq 1$, let

$$f_\mu(z) = z^3 - \frac{2\xi}{\sqrt{2(1-\zeta)}}z^2 - \frac{1}{4}\left(2\zeta - 6 + \frac{2\xi^2}{\zeta - 1}\right)z.$$

(Note that for $\mu = 0$, f_μ is equal to the previously defined f_0.) Then the fixed points of f_μ are 0, $\alpha = \frac{\xi}{\sqrt{2(1-\zeta)}} + \sqrt{\frac{1-\zeta}{2}}\,i$ and $\bar{\alpha}$ and the multipliers of α and $\bar{\alpha}$ are μ and $\bar{\mu}$ respectively. Hence if $|\mu| < 1$, then $f_\mu \in \mathcal{C}(\lambda_0)$. Fix $k > 1/2$ and let $\mu_n = 1 - k/n^2 + i/n$. Then $|\mu_n| < 1$ for sufficiently large n, $\mu_n \to 1$, and



FIGURE 10. The Julia sets of f_{μ_n} and its limit.

$$f_{\mu_n}(z) \to g_k(z) = z^3 - \sqrt{\frac{2}{k}}z^2 + \left(1 + \frac{1}{2k}\right)z.$$

Clearly, g_k is real and has a repelling fixed point at 0 and a parabolic fixed point at $1/\sqrt{2k} > 0$, hence $K(g_k) \cap \mathbb{R}$ is a closed interval and the external rays of angles 0 and $1/2$ do not land at the same point, hence $g_k \notin \mathcal{C}(\lambda_0)$ (see Figure 10). In particular, $\mathcal{C}(\lambda_0)$ is not compact.

Another example is given in Figure 11, which was proposed by Goldberg-Milnor [GM].



FIGURE 11. Butterfly collapses to a period two parabolic cycle. See Goldberg-Milnor [GM].

8. SURJECTIVITY OF STRAIGHTENING MAPS

8.1. **Preliminaries.** To prove our surjectivity results we will employ the compactness of primitive renormalizable polynomials and the injectivity of straightening proven in previous sections together with the nice properties of straightening of quadratic-like families, established by Douady and Hubbard in [DH2]. We summarize some of these properties as follows (compare with [DH2, Proposition 13 and Chapter IV]).

Theorem 8.1. *Let X be a complex manifold and $\{f_\mu\}_{\mu\in X}$ be a holomorphic family of quadratic-like maps. Let*

$$\mathcal{M}_X = \{\mu \in X; K(f_\mu) \text{ is connected }\},$$

and denote by

$$\chi : \mathcal{M}_X \to \mathcal{M}$$

the corresponding straightening map. That is, $\chi(\mu) = c$ where $c \in \mathcal{M}$ is such that $z \mapsto z^2 + c$ is hybrid equivalent to f_μ. Then the following statements hold:

- *χ is continuous.*
- *For all $c \in \mathcal{M}$, we have that $\chi^{-1}(c)$ is a complex analytic subspace of X.*
- *If X has complex dimension one and $\mu_0 \in \mathcal{M}_X$, then there exists a neighborhood V of μ_0 in $\mathcal{M}_X$ such that χ is either constant on V or $\chi(V)$ contains a neighborhood of μ_0 in $\mathcal{M}$.*

We will also need the following lemma on quasiconformal deformations:

Lemma 8.2. *Let λ_0 be a d-invariant rational lamination. Let $f \in \mathcal{R}(\lambda_0)$ and $\chi_{\lambda_0}(f) = P \in C(T(\lambda_0))$. Consider a quasiconformal deformation of P on $K(P)$, i.e., there exists a quasiconformal map $\phi : \mathbb{C} \to \mathbb{C}$ whose complex dilatation is supported on $K(P)$ such that $\phi \circ P = P_1 \circ \phi$ for some $P_1 \in C(T(\lambda_0))$.*

Then there exists a quasiconformal deformation $f_1 \in \mathcal{R}(\lambda_0)$ of f on $K(f)$ such that $\chi_{\lambda_0}(f_1) = P_1$.

Proof. Let us denote by σ_0 the standard complex structure on $\mathbb{C}$ and let $\sigma = \phi^*\sigma_0$. Define an almost complex structure σ_1 on $\mathbb{C}$ as follows:

$$\sigma_1 = \begin{cases} (\psi_v \circ f^n)^*\sigma & \text{on } f^{-n}(K_f(v)) \text{ for some } v \in |T(\lambda_0)| \text{ and } n \geq 0, \\ \sigma_0 & \text{otherwise,} \end{cases}$$

where $\psi = (\psi_v)$ is a hybrid conjugacy from a λ-renormalization of f to P. Then σ is well-defined because σ is g-invariant. Its dilatation is uniformly bounded because σ is defined by pullbacks of $\psi_v^*\sigma = (\phi \circ \psi_v)^*\sigma_0$ by holomorphic maps and pullbacking by holomorphic map does not change the maximal dilatation. Furthermore, σ is f-invariant by construction. Therefore, by the measurable Riemann mapping theorem, $f : (\mathbb{C}, \sigma_1) \to (\mathbb{C}, \sigma_1)$ is conformally conjugate to a polynomial $f_1 : \mathbb{C} \to \mathbb{C}$. Since this deformation preserves external rays and their landing points, we have $f_1 \in C(\lambda_0)$ and λ_0-renormalization of f gives the λ_0-renormalization of f_1. More precisely, let $\phi_1 : \mathbb{C} \to \mathbb{C}$ be the quasiconformal homeomorphism such that $\phi_1^*\sigma_0 = \sigma_1$ and $f_1 = \phi_1 \circ f \circ \phi_1^{-1}$. Take a λ_0-renormalization $g = (f : U'_v \to U_{\sigma_{\lambda_0}(v)})$ of f. Then $g_1 = (f_1 : \phi_1(U'_v) \to \phi_1(U_{\sigma_{\lambda_0}(v)}))$ is a λ_0-renormalization of f_1 and it is hybrid equivalent to P_1 by $\phi \circ \psi \circ \phi_1^{-1}$. □

8.2. **Surjectivity of straightening: primitive disjoint type.**

Theorem 8.3. *Let $f_0 \in C(d)$ be an internally angled primitive hyperbolic post-critically finite polynomial with rational lamination λ_0. If f_0 has exactly $d-1$ superattracting periodic orbits, then the corresponding straightening map $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \to \mathcal{M}^{d-1}$ is a homeomorphism.*

Proof. Recall that the interior of the Mandelbrot set $\mathcal{M}$ is dense in $\mathcal{M}$. Therefore, the interior of $\mathcal{M}^{d-1}$ is also dense in $\mathcal{M}^{d-1}$. The components of the interior of $\mathcal{M}$ are either hyperbolic (maps with an attracting cycle) or queer. The closure of the union of the hyperbolic components is the complement of the queer components. That is, $\partial\mathcal{M}$ is contained in the closure of the union of the hyperbolic components.

Let λ_0 be an internally angled hyperbolic primitive rational lamination of degree d with reduced mapping schema of disjoint type. Given $d-1$ components $W_1, \ldots, W_{d-1}$ of the interior of $\mathcal{M}$ we will show that $S_0 = W_1 \times \cdots \times W_{d-1}$ is in the image of the associated straightening map χ_{λ_0}. Since $\mathcal{R}(\lambda_0)$ is compact and χ_{λ_0} is continuous, this implies that χ_{λ_0} is surjective.

Since hyperbolic maps are in the image of χ_{λ_0}, we have that $(\partial\mathcal{M})^{d-1}$ is also contained in the image. In particular, $\partial W_1 \times \cdots \times \partial W_{d-1} \subset \chi_{\lambda_0}(\mathcal{R}(\lambda_0))$. Now, for $n \in \{1, \ldots, d-1\}$, let

$$S_n = \partial W_1 \times \cdots \partial W_n \times W_{n+1} \times \cdots \times W_{d-1}.$$

Observe that $S_{d-1} \subset \chi_{\lambda_0}(\mathcal{R}(\lambda_0))$. To prove that S_0 is contained in $\chi_{\lambda_0}(\mathcal{R}(\lambda_0))$ it is enough to establish the following.

Claim. For all $1 \leq n \leq d-1$, if $S_n \subset \chi_{\lambda_0}(\mathcal{R}(\lambda_0))$, then $S_{n-1} \subset \chi_{\lambda_0}(\mathcal{R}(\lambda_0))$.

Proof of the claim. Let $\mathbf{c} = (c_1, \ldots, c_d) \in S_n$. Denote by $\pi : \mathbb{C}^{d-1} \to \mathbb{C}^{d-2}$ the projection which forgets the nth coordinate and by $\pi_n : \mathbb{C}^{d-1} \to \mathbb{C}$ the projection onto the nth coordinate . By Theorem 8.1, $(\pi \circ \chi_{\lambda_0})^{-1}(\mathbf{c})$ is a complex analytic space Y, since post-composition of χ_{λ_0} by projection onto one coordinate is the straightening map of a quadratic-like family. After resolution of singularities, we may assume that Y is a complex manifold. Moreover, $\pi_n \circ \chi_{\lambda_0} : \mathcal{R}(\lambda_0) \cap Y \to \mathcal{M}$ is injective, by Theorem B. Since $\mathcal{R}(\lambda_0)$ is compact (Theorem D) and the uncountable set ∂W_n is contained in $\pi_n \circ \chi_{\lambda_0}(\mathcal{R}(\lambda_0) \cap Y)$, there exists $f \in \mathcal{R}(\lambda_0)$ contained in a component of dimension at least one of Y such that $\pi_n \circ \chi_{\lambda_0}(f) \in \partial W_n$. Theorem B implies that $\pi_n \circ \chi_{\lambda_0}$ is not constant along any one dimensional submanifold of Y containing f, therefore $\pi_n \circ \chi_{\lambda_0}(\mathcal{R}(\lambda_0) \cap Y)$ contains a neighborhood of c_n in $\mathcal{M}$, by Theorem 8.1. From Lemma 8.2, we conclude that $\pi_n \circ \chi_{\lambda_0}(\mathcal{R}(\lambda_0) \cap Y)$ contains W_n. Hence, for all $\mathbf{c} = (c_1, \dots, c_d) \in S_n$ we have that $\{c_1\} \times \cdots \times \{c_{n-1}\} \times W_n \times \{c_{c+1}\} \times \cdots \times \{c_{d-1}\} \subset \chi_{\lambda_0}(\mathcal{R}(\lambda_0))$ and the claim follows. ◻

8.3. **Surjectivity of straightening: cubic primitive capture type.**

Theorem 8.4. *Let $f_0 \in \mathcal{C}(3)$ be an internally angled primitive hyperbolic post-critically finite polynomial with rational lamination λ_0 and reduced mapping schema of capture type. then the associated straightening map $\chi_{\lambda_0} : \mathcal{R}(\lambda_0) \to \mathcal{MK}$ is a bijection.*

The proof of Theorem 8.4 is similar to the disjoint case. However we have to be careful since the straightening maps involved are discontinuous [In4]. Nevertheless, we use the fact that these maps are continuous along carefully chosen sequences.

Lemma 8.5. *Let $f_n : U_n' \to U_n$ be a sequence of quadratic-like maps with connected Julia sets. Assume that the following hold:*

(i) *f_n converges to a quadratic-like map $f : U' \to U$. More precisely, $f_n \to f$ uniformly on some neighborhood of $K(f; U', U)$ as $n \to \infty$;*
(ii) *f_n is hybrid equivalent to $g_n \in \mathcal{M}$.*
(iii) *$g_n \to g \in \mathcal{M}$;*
(iv) *For any $z \in \operatorname{int} K(g)$, $z \in \operatorname{int} K(g_n)$ for all sufficiently large n.*

Then we can choose hybrid conjugacy ψ_n between f_n and g_n such that ψ_n converges to a hybrid conjugacy ψ between f and g.

In particular, let λ_0 be a hyperbolic 3-invariant rational lamination of capture type. Let $f_n \to f$ be a convergent sequence in $\mathcal{R}(\lambda_0)$. If the quadratic renormalization $f_n^p : U_n' \to U_n$ of f_n satisfies the above assumption, then $\chi_{\lambda_0}(f_n) \to \chi_{\lambda_0}(f)$.

Remark 8.6. The assumption (iv) is equivalent to $J(g_n) \to J(g)$ in the Hausdorff topology. Furthermore, (iv) holds if one of the following hold:

(i) $J(g) = K(g)$, i.e., $\operatorname{int} K(g)$ is empty;
(ii) g lies in the boundary of a hyperbolic component $\mathcal{W}$ of $\mathcal{M}$ and $g_n \in \mathcal{W} \to g$ non-tangentially (see [Mc2] for the case g parabolic and see [Yo] for the case g has a Siegel disk).

Proof. Using the tubing construction of hybrid conjugacies [DH2], we can always choose hybrid conjugacies ψ_n so that there exists some $K \geq 1$ such that ψ_n is K-quasiconformal for any n. Therefore, by passing to a subsequence, ψ_n converges to some quasiconformal map ψ. Furthermore, they also proved that hybrid conjugacy constructed by the tubing construction depends continuously outside the boundary of $\mathcal{M}$.

Hence we may assume $g \in \partial\mathcal{M}$. By Theorem 8.1, f is hybrid equivalent to g and ψ is a quasiconformal conjugacy between f and g. Let $\mu = \frac{\bar\partial\psi}{\partial\psi}$ be the complex dilatation of ψ. Then, by the equation $\psi \circ f = g \circ \psi$, we have $f^*\mu = \mu$. Since g does not carry an invariant line field on its Julia set (otherwise, g must lie in a queer component), we may assume that

μ vanishes on $J(g)$. Furthermore, since ψ_n is holomorphic in the interior of $K(f_n)$, μ also vanishes on $\operatorname{int} K(g)$ by (iv). Therefore, ψ is a hybrid conjugacy between f and g.

For $f_n \to f \in \mathcal{R}(\lambda_0)$, let ω_n be the captured critical point for f_n and k be the capture time. Then we have $\chi_{\lambda_0}(f_n) = (g_n, x_n)$ where $x_n = \psi_n(f_n^k(\omega_n))$. If $f_n : U_n' \to U_n$ satisfy the assumption, then, after passing to the limit, we have $x = \psi(f^k(\omega))$, where ω is the captured critical point for f. Therefore, $\chi_{\lambda_0}(f) = (g, x) = \lim \chi_{\lambda_0}(f_n)$. ◻

Proof of Theorem 8.4. To simplify notation, here we identify the map $h(z) = z^2 + c$ with $c \in \mathbb{C}$. That is, straightening takes values in the set:

$$\mathcal{MK} = \{(h, x);\ h(z) = z^2 + c \in \mathcal{M},\ x \in K(h)\}$$

Observe that if h is hyperbolic and $x \in \operatorname{int} K(h_0)$, then the pair (h, x) represents a hyperbolic dynamical system over the reduced schema of λ_0. By Theorem C, (h, x) is in the image of χ_{λ_0}.

Now we consider $(h_0, x_0) \in \mathcal{MK}$. To prove that that $(h_0, x_0) \in \mathcal{R}(\lambda_0)$ we consider two cases according to whether h_0 is in a queer component or not.

Case I: h_0 does not lie in any queer component.

Take sequences $h_n \to h_0$ and $x_n \to x_0$ satisfying (iv) in Lemma 8.5 such that h_n is hyperbolic and $x_n \in \operatorname{int} K(h_n)$. Let $f_n \in \mathcal{R}(\lambda_0)$ satisfy $\chi_{\lambda_0}(f_n) = (h_n, x_n)$. Then, by Lemma 8.5, we have $\chi_{\lambda_0}(f) = (h_0, x_0)$.

When h_0 is in a queer component we further subdivide into two cases according to whether the captured other critical forward orbit is finite or not.

Case II.a: Assume that $h_0 \in \mathcal{W}$ for some queer component $\mathcal{W}$ and that x_0 is eventually periodic for h_0.

In this case, we have $h_0^j(x_0) = h_0^k(x_0)$ for some $j \neq k$. Take $h_1 \in \partial\mathcal{W} \subset \mathcal{M}$ and an eventually periodic point x_1 for h_1, which is the continuation of x_0 along a path in $\overline{\mathcal{W}}$. Take $f_1 \in \mathcal{R}(\lambda_0)$ with $\chi_{\lambda_0}(f_1) = (h_1, x_1)$. Let ω_1 be the captured critical point for f_1 and let $l > 0$ be the capture time. Then we have $f_1^{j+l}(\omega_1) = f_1^{k+l}(\omega_1)$.

Now consider the one-dimensional complex analytic sets

$$V = \{(f, \omega);\ f \in \mathrm{Poly}(3),\ \omega\text{: critical point for } f \text{ and } f^{j+l}(\omega) = f^{k+l}(\omega)\},$$

$$\tilde{V} = \{(h, x);\ h \in \mathrm{Poly}(2),\ h^j(x) = h^k(x)\}.$$

Then $(f_1, \omega_1) \in V$ and $\chi_{\lambda_0}(f_1) = (h_1, x_1)$ lies in $\tilde{V}$. Let $\tilde{\chi}(f, \omega) = \pi_0 \circ \chi_{\lambda_0}(f)$. By Theorem 8.1, we can extend $\tilde{\chi}$ continuously in a small neighborhood $\mathcal{U}$ of f_1 in V. If $\tilde{\chi}$ is constant on $\mathcal{U}$, then χ_{λ_0} is also constant and it contradicts the injectivity of χ_{λ_0}. Hence $\tilde{\chi}$ is non-constant, and we can find $(f_2, \omega_2) \in \mathcal{U}$ with $\tilde{\chi}(f_2, \omega_2) \in \mathcal{W}$. Furthermore, if we move f continuously from f_1 to f_2 in $\tilde{\chi}^{-1}(\mathcal{W}) \cup \{f_1\}$, then every periodic point does not bifurcate, since they are all repelling for parameters in the closure of $\tilde{\chi}^{-1}(\mathcal{W})$. This implies that ω_2 is still a captured critical point. Therefore, $f_2 \in \mathcal{R}(\lambda_0)$ and $\chi_{\lambda_0}(f_2) \in \tilde{V}$. As before, deforming f_2 quasiconformally, we obtain a polynomial $f \in \mathcal{R}(\lambda_0)$ with $\chi_{\lambda_0}(f) = (h_0, x_0)$.

Case II.b: Assume that $h_0 \in \mathcal{W}$ for some queer component $\mathcal{W}$ and $x_0 \in J(h_0) (= K(h_0))$.

Since periodic points are dense in $J(h_0)$, we can take a sequence $x_n \to x_0$ with x_n periodic. Take $f_n \in \mathcal{R}(\lambda_0)$ which satisfy $\chi_{\lambda_0}(f_n) = (h_0, x_n)$. Since $\mathcal{R}(\lambda_0)$ is compact, we may assume f_n converges to some $f \in \mathcal{R}(\lambda_0)$. By Lemma 8.5, we have $\chi_{\lambda_0}(f) = (h_0, x_0)$. ◻

8.4. **Complex submanifolds and quasiconformality of straightening.** Here we prove some regularity properties of straightening maps of primitive disjoint type and primitive cubic capture type. That is, we finish the proof of theorems E and F.

8.4.1. *Proof of Theorem E.* In view of Theorem 8.3 the straightening map χ_{λ_0} is surjective. We must show that for any $0 \leq k \leq d-1$, given $1 \leq i_1 < \cdots < i_k \leq d-1$ and $c_1, \ldots, c_k \in \mathcal{M}$, the set

$$\{f \in \mathcal{R}(\lambda_0); \chi_{i_j}(f) = c_j \text{ for all } j = 1, \ldots, k\}$$

is contained in a codimension k complex submanifold $\mathcal{S}$ of Poly(d). For $k = 0$, then the statement above is clearly true. We proceed recursively and assume that we have proven the statement for $k \geq 0$. So let $\mathcal{S}$ be a codimension k complex submanifold such that for all $f \in \mathcal{R}(\lambda_0) \cap \mathcal{S}$ and for all $j = 1, \ldots, k$, we have that $\chi_{i_j(f)}(f) = c_j$. We work in the space $\mathcal{QL}$ of quadratic-like germs introduced by [L]. Let

$$\iota(f) = [f^\ell : U' \to U] \in \mathcal{QL}$$

be the quadratic-like map obtained from $f \in \mathcal{U}$ which is hybrid conjugate to $z^2 + \chi_{i_{k+1}}(f)$. Since ι is an analytic map and by [L, Lemma 4.25], $\iota(\mathcal{S})$ is transverse to the foliation $\mathcal{F}$ of hybrid classes at all $f \in \mathcal{S} \cap \mathcal{R}(\lambda_0)$. Therefore, the subset of formed by polynomial $f \in \mathcal{S}$ such that $\iota(f)$ is hybrid conjugate to $z \mapsto z^2 + c_{i_{k+1}}$ is a codimension one submanifold of $\mathcal{S}$.

The quasiconformality of χ along one dimensional slices as in the statement of Theorem E, follows directly from [L, Theorem 4.26].

8.4.2. *Capture case.* In order to complete the proof of Theorem F we must establish that the straightening map involved is quasiconformal along appropriate slices. The proof is similar than that of Buff and Henriksen in [BHe]: We construct a holomorphic motion on the filled Julia set and use its quasiconformal extension, obtained via the λ-lemma, to show that the straightening map has the desired quasiconformal regularity.

Proof of Theorem F. Let $c \in \mathcal{M}$ and $\mathcal{S}_c$ be as in the statement of the theorem. From Douady-Hubbard's theorem 8.1 it follows that $\mathcal{S}_c$ is a complex analytic subspace of Poly(3). We take a point $f_0 \in \mathcal{R}(\lambda_0) \cap \mathcal{S}_c$ and denote by $\mathcal{S}$ a branch of $\mathcal{S}_c$ at f_0. We will show that χ is quasiconformal on $\mathcal{S}$. Then, by injectivity, we will conclude that there is at most one branch of $\mathcal{S}_c$ at f_0. Hence, it will immediately follow that $\mathcal{S}_c$ is locally irreducible.

Consider a family $(f^{\ell_{v_0}} : U'_{v_0} \to U_{v_0})_{f \in \mathcal{S}}$ of quadratic-like mappings. Since the hybrid class of the elements of this family is constant, it is stable on the whole parameter space $\mathcal{S}$, in the sense of Mañé-Sad-Sullivan. It follows that the Julia set $J_f(v_0) = \partial K_f(v_0)$ admits a holomorphic motion on $\mathcal{S}$ [Mc1]. Furthermore, we have a natural conformal conjugacy $\psi_{f_0}^{-1} \circ \psi_f : \operatorname{int} K_{f_0}(v_0) \to \operatorname{int} K_f(v_0)$ between f_0 and f, which is also a holomorphic motion on $\mathcal{S}$, where ψ_f is a hybrid conjugacy between $f^{\ell_{v_0}} : U'_{v_0} \to U_{v_0}$ and $Q(z) = z^2 + c$. Gluing both holomorphic motions together, we obtain a holomorphic motion $h_f : K_{f_0}(v_0) \to K_f(v_0)$. For $f \in \mathcal{S} \cap \mathcal{R}(\lambda_0)$, we have

$$\chi(f) = (c, \psi_f(f^{\ell_{v_1}}(\omega_f))) = (c, \psi_{f_0} \circ h_f(f^{\ell_{v_1}}(\omega_f))),$$

where ω_f is the captured critical point.

By applying the λ-lemma [MSS] [Sl], h_f extends to a holomorphic motion

$$h : \mathcal{S} \times \mathbb{C} \to \mathbb{C},$$

such that $h_f : \mathbb{C} \to \mathbb{C}$ is quasiconformal. By the same argument as [BHe, Lemma 13], the map

$$f \mapsto h_f(f^{\ell_{v_1}(\omega_f)})$$

is locally quasiconformal on $\mathcal{S}$. Taking a finite covering of $\mathcal{R}(\lambda_0) \cap \mathcal{S}_c$, we conclude that $\chi : \mathcal{R}(\lambda_0) \cap \mathcal{S}_c \to K(Q)$ extends to a quasiconformal homeomorphism.

To finish the proof we have to show that $\mathcal{R}(\lambda_0)$ is connected. In fact, observe that if C is both open and closed in $\mathcal{R}(\lambda_0)$ and $\mathcal{S}_c$ is as above, then $\mathcal{S}_c$ is either contained in C or disjoint from C. Now $\chi_1 : \mathcal{R}(\lambda_0) \to \mathcal{M}$ is continuous and C is compact. Therefore, $\chi_1(C)$ and $\chi_1(\mathcal{R}(\lambda_0) \setminus C)$ are closed and disjoint subsets of $\mathcal{M}$. Then one of these sets must be empty, since $\mathcal{M}$ is connected. Hence, either C is empty or $C = \mathcal{R}(\lambda_0)$. $\square$

References

[BrH] B. Branner and J.H. Hubbard, *The iteration of cubic polynomials. I. The global topology of parameter space.*, Acta Math. **160** (1988), 143–206.

[BHe] Buff, X. and Henriksen, Christian, *Julia Sets in Parameter Spaces*, Comm. Math. Phys. **220** (2001), 333–375.

[DH1] A. Douady, J. H. Hubbard, *Étude dynamique des polynômes complexes* I & II. Publ. Math. d'Orsay (1984-1985).

[DH2] A. Douady, J. H. Hubbard, *On the dynamics of polynomial-like mappings.* Ann. Sci. École Norm. Sup. (4) **18** (1985), 287–343.

[EY] A. Epstein, M. Yampolsky, *Geography of the cubic connectedness locus: intertwining surgery.* Ann. Sci. École Norm. Sup. (4) **32**

[GM] L. Goldberg, J. Milnor. *Fixed points of polynomial maps. II. Fixed point portraits.* Ann. Sci. École Norm. Sup. (4) **26** (1993), 51–98.

[H] J. H. Hubbard, *Local connectivity of Julia sets and bifurcation loci: three theorems of J.-C. Yoccoz.* Topological methods in modern mathematics, 467–511, Publish or Perish, Houston, TX, 1993.

[Ha1] P. Haïssinsky, *Chirurgie parabolique.* C. R. Acad. Sci. Paris Sér. I Math. **327** (1998), no. 2, 195–198.

[Ha2] P. Haïssinsky, *Modulation dans l'ensemble de Mandelbrot.* The Mandelbrot set, theme and variations, 37–65, London Math. Soc. Lecture Note Ser., 274, Cambridge Univ. Press, Cambridge, 2000.

[Ha3] P. Haïssinsky, *Chirurgie croisée*. Bull. Soc. mathḞrance, **128** (2000), 599–654.

[Ha4] P. Haïssinsky, *Pincement de polynômes*. Comment. Math. Helv. 77 (2002), 1–23.

[He] C. Henriksen, *The combinatorial rigidity conjecture is false for cubic polynomials*, Trans. Amer. Math. Soc. **355** (2003), 3625–3639.

[In1] H. Inou, *Renormalization and rigidity of polynomials of higher degree.* J. Math. Kyoto Univ. **42** (2002), 351–392.

[In2] H. Inou, *Homeomorphisms from Julia sets into connectedness loci.* Ergodic Theory & Dynamical Systems **26** (2006), 1087–1113.

[In3] H. Inou, *Limits of renormalizable polynomials*. Nonlinearity **19** (2006), 1769–1799.

[In4] H. Inou, *Combinatorics and topology of straightening maps II: discontinuity*. In preparation.

[Ki] J. Kiwi, *Rational laminations of complex polynomials*. Laminations and foliations in dynamics, geometry and topology (Stony Brook, NY, 1998), 111–154, Contemp. Math., 269, Amer. Math. Soc., Providence, RI, 2001.

[L] M. Lyubich, *Feigenbaum-Coullet-Tresser universality and Milnor's hairiness conjecture*. Ann. of Math. (2) **149** (1999), 319–420.

[MSS] Mañé, R.; Sad, P.; Sullivan, D. *On the dynamics of rational maps.* Ann. Sci. École Norm. Sup. (4) **16** (1983), p. 193-217.

[Mc1] C. T. McMullen, *Complex dynamics and renormalization.* Annals of Mathematics Studies, 135. Princeton University Press, Princeton, NJ, 1994.

[Mc2] C. T. McMullen, *Hausdorff dimension and conformal dynamics. II. Geometrically finite rational maps.* Comment. Math. Helv. **75** (2000), 535–593.

[Mc3] C. T. McMullen, *The Mandelbrot set is universal.* The Mandelbrot set, theme and variations, 1–17, London Math. Soc. Lecture Note Ser., 274, Cambridge Univ. Press, Cambridge, 2000.

[MS] C. T. McMullen, D. P. Sullivan, *Quasiconformal homeomorphisms and dynamics. III. The Teichmüller space of a holomorphic dynamical system.* Adv. Math. **135** (1998), 351–395.

[Mi1] J. Milnor, *Remarks on iterated cubic maps.* Experiment. Math. **1** (1992), 5–24.

[Mi2] J. Milnor, *Local Connectivity of Julia Sets: Expository Lectures*. The Mandelbrot set, theme and variations, 67–116, London Math. Soc. Lecture Note Ser., 274, Cambridge Univ. Press, Cambridge, 2000.

[Mi3] J. Milnor, *Dynamics in one complex variable. Third edition.* Annals of Mathematics Studies, 160. Princeton University Press, Princeton, NJ, 2006.

[Mi4] J. Milnor, *Hyperbolic components in spaces of polynomial maps*. Stony Brook IMS preprint 1992/3.

[Po] A. Poirier, *On postcritically finite polynomials, part one: critical portraits*. Stony Brook IMS preprint 1993/5.

[Sch] D. Schleicher, *On fibers and local connectivity of Mandelbrot and Multibrot sets*. Fractal geometry and applications: a jubilee of Benoît Mandelbrot. Part 1, Proc. Sympos. Pure Math. 72 (2004), 477–517.

[Sl] Z. Slodkowski, *Extensions of holomorphic motions.* Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) 22 (1995) 185–210.
[Th] W. Thurston, *On the combinatorics of iterated rational maps*, manuscript, 1985.
[Yo] J. C. Yoccoz, *Théorème de Siegel, nombres de Bruno et polynômes quadratiques*. Petits diviseurs en dimension 1. Astérisque No. 231 (1995), 3–88.

DEPARTMENT OF MATHEMATICS, KYOTO UNIVERSITY, KYOTO 606-8502, JAPAN.
URL: `www.math.kyoto-u.ac.jp/˜inou/`

FACULTAD DE MATEMÁTICAS, PONTIFICIA UNIVERSIDAD CATÓLICA DE CHILE.
URL: `www.mat.puc.cl/˜jkiwi`